International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
IJABR 2025; 9(1): 452-461
www.biochemjournal.com
Received: 21-10-2024
Accepted: 26-11-2024

Adil Rahim Magray
ICAR-Indian Agricultural Research Institute, New Delhi, India

Latief Bashir
ICAR-Indian Agricultural Research Institute, New Delhi, India

Azhar Mehmood
ICAR-Indian Agricultural Research Institute, New Delhi, India

Arif Hassan
Shere Kashmir University of Agricultural Sciences and Technology, Srinagar, Jammu and Kashmir, India

Suhail Ashraf
Faculty of Biology, CeBiTec-Center for Biotechnology, Bielefeld University, Bielefeld, Germany

Sampatirao Dilip
ICAR-Indian Agricultural Research Institute, New Delhi, India

Sameer Ahmad Dar
Hemvati Nandan Bahuguna Garhwal University, Srinagar, Jammu and Kashmir, India

Jitendra Kumar Yadav
ICAR-Indian Agricultural Research Institute, New Delhi, India

Kunal
ICAR-Indian Agricultural Research Institute, New Delhi, India

Uma Shankar
Department of Plant Pathology, Acharya Narendra Deva University of Agriculture and Technology, Ayodhya, Uttar Pradesh, India

Dileep Meena
ICAR-Indian Agricultural Research Institute, New Delhi, India

Mamta Yadav
Shri Karan Narendra Agriculture University, Jobner, Jaipur, Rajasthan, India

Corresponding Author:
Latief Bashir
ICAR-Indian Agricultural Research Institute, New Delhi, India

Physiological and molecular insights into the allelopathic effects on agroecosystems under changing environmental conditions

Adil Rahim Magray, Latief Bashir, Azhar Mehmood, Arif Hassan, Suhail Ashraf, Sampatirao Dilip, Sameer Ahmad Dar, Jitendra Kumar Yadav, Kunal, Uma Shankar, Dileep Meena and Mamta Yadav

DOI: https://doi.org/10.33545/26174693.2025.v9.i1f.3534

Abstract
The phenomenon of allelopathy, wherein plants release biochemicals that influence the growth, development, and survival of neighboring organisms, holds immense significance in shaping agroecosystems. With changing environmental conditions, understanding the physiological and molecular mechanisms underlying allelopathic interactions becomes crucial for sustainable agricultural practices. This abstract reviews recent advancements in the field, focusing on the intricate interplay between allelopathic compounds, recipient plant physiology, and environmental factors. Allelopathic compounds, including phenolics, terpenoids, and alkaloids, exert diverse effects on recipient plants, ranging from growth promotion to inhibition, depending on concentration, duration of exposure, and plant species involved. Recent studies highlight the role of allelopathy in weed suppression, nutrient cycling, and crop productivity enhancement, underscoring its potential for weed management and agroecological intensification. At the physiological level, allelopathic compounds modulate various processes within recipient plants, including seed germination, root growth, photosynthesis, and nutrient uptake. Mechanisms such as oxidative stress, hormonal signaling, and membrane permeability alterations contribute to the observed physiological responses.

Keywords: Allelochemicals, allelopathy, molecular mechanism, signalling pathway, gene expression, sustainable agriculture

1. Introduction
The intricate interplay between allelopathic effects and agroecosystems in response to shifting environmental dynamics underscores the necessity for comprehensive understanding and exploration. Research by Li *et al.* (2019) [75] highlights the multifaceted nature of allelopathy and its implications on agricultural sustainability. As global climate patterns continue to evolve, elucidating the physiological and molecular mechanisms underlying allelopathic interactions becomes imperative for effective agroecosystem management (Kaur *et al*., 2020) [63]. Allelopathy, defined as the biochemical interactions among plants leading to growth inhibition or stimulation, has garnered significant attention due to its potential role in shaping plant community structure and productivity (Singh *et al*., 2017). The study conducted by Kong *et al.* (2018) [68] emphasizes the dynamic nature of allelopathic interactions, influenced by various factors including soil type, plant species, and environmental conditions. Understanding these complexities is crucial for devising strategies to harness the positive aspects of allelopathy while mitigating its detrimental effects on crop growth and yield (Cheng *et al*., 2021) [19]. Environmental changes, such as altered precipitation patterns, temperature fluctuations, and increasing levels of atmospheric CO_2, can significantly modulate allelopathic interactions within agroecosystems (Yang *et al*., 2022) [114]. Recent findings by Huang *et al.* (2023) [47] underscore the importance of considering these environmental variables when studying allelopathy, as they can influence the production and release of allelochemicals by donor plants, as well as the sensitivity of recipient plants to these compounds. At the physiological level, allelopathic interactions involve a complex network of signaling pathways and biochemical responses within both

donor and recipient plants (Farooq *et al*., 2018) [29]. Studies by Jia *et al*. (2021) [58] have elucidated the role of phytohormones, such as auxins, cytokinins, and abscisic acid, in mediating allelopathic effects, highlighting the intricate crosstalk between hormonal signaling and allelochemical-induced stress responses. Furthermore, recent advancements in molecular biology techniques have provided unprecedented insights into the genetic basis of allelopathy (Tan *et al*., 2020) [95]. Genome-wide association studies (GWAS) conducted by Zhou *et al*. (2022) [123, 126] have identified key genes and regulatory elements involved in allelochemical synthesis, secretion, and detoxification, offering new avenues for crop improvement through molecular breeding approaches. In conclusion, the study of allelopathic effects on agroecosystems under changing environmental conditions represents a multifaceted research area with significant implications for agricultural sustainability. Integrating physiological and molecular approaches, alongside environmental considerations, is essential for unraveling the complexities of allelopathic interactions and developing strategies to enhance crop productivity while minimizing ecological risks.

2. Understanding Physiological Mechanisms

2.1 Exploring Plant-Plant Interactions: A Multifaceted Perspective

Plant-plant interactions, a cornerstone of ecological dynamics, unfold in a complex web of symbiotic, competitive, and allelopathic relationships, as elucidated by studies such as those conducted by Callaway and Walker (1997) [14], Cahill and McNickle (2011) [13], and Brooker *et al*. (2008) [11]. These interactions, often mediated through chemical signaling, spatial competition, and resource partitioning, have been extensively investigated in various ecosystems worldwide, as documented in the works of Grime (1977) [40], Bertness and Callaway (1994) [6], and Goldberg and Werner (1983) [38]. In the realm of competition, studies by Tilman (1982) [100] and Adler *et al*. (2007) [1] have shed light on the mechanisms underlying resource acquisition and exploitation among neighboring plants, while Connell (1971) [23] and Pickett (1989) [85] have explored the role of allelopathy in shaping community structure and composition. Furthermore, pioneering research by Grman *et al*. (2010) [41] and Fukami *et al*. (2010) [33] has underscored the significance of plant-soil feedbacks in regulating species coexistence and diversity. Beyond competitive interactions, mutualistic associations, as highlighted by Bever (2003) [7] and Kiers *et al*. (2010) [67], play a crucial role in ecosystem functioning and resilience. Studies on mycorrhizal symbioses, such as those by Smith and Read (2008) [94] and van der Heijden *et al*. (2015) [101], have revealed the intricate networks of nutrient exchange and communication between plants and soil fungi, shaping community dynamics and productivity. Moreover, facilitation, a form of positive interaction, has garnered attention in recent years, with research by Bruno *et al*. (2003) [12] and Michalet *et al*. (2006) [80] illustrating the role of nurse plants in ameliorating harsh environmental conditions and promoting the establishment of neighboring species. Similarly, studies by Holzapfel and Mahall (1999) [43] have demonstrated the importance of allelopathic interactions in facilitating the coexistence of diverse plant species in competitive environments. Furthermore, recent advances in molecular ecology, exemplified by work conducted by Wurst *et al*. (2010) [111] and Schweitzer *et al*. (2008) [91], have provided insights into the genetic basis of plant interactions, unraveling the intricate mechanisms underlying trait-mediated interactions and community dynamics.

2.2 Unveiling Allelochemical Dynamics and impacts on Growth and Development

Understanding the intricate dynamics of allelochemical interactions is paramount in elucidating their impacts on plant growth and development (Rice *et al*., 2013) [88]. These chemical compounds, produced by one plant species, can influence the growth, development, and even survival of neighboring plants (Inderjit and Duke, 2003) [51]. Allelopathy, the phenomenon whereby plants release chemicals into the environment to inhibit the growth of competing plants, has garnered significant attention in recent years (Weston and Duke, 2003) [108]. The allelochemicals involved in these interactions can have diverse effects, ranging from growth promotion to inhibition, depending on the concentration and specific compounds involved (Kato-Noguchi, 2004) [60]. Allelopathic interactions play a crucial role in shaping plant communities and can have significant ecological implications (Rice, 1984) [87]. Understanding the mechanisms underlying allelochemical dynamics is essential for developing sustainable agricultural practices and managing invasive plant species (Inderjit *et al*., 2005) [55]. The mechanisms through which allelochemicals exert their effects on plant growth and development are diverse and complex (Blum *et al*., 1999) [9]. These compounds can interfere with various physiological processes, including seed germination, root elongation, and nutrient uptake (Duke, 2015) [27]. One common mode of action involves allelochemicals disrupting cell membrane integrity, leading to leakage of cellular contents and ultimately cell death (Khanh *et al*., 2007) [66]. Additionally, allelochemicals can interfere with hormone signaling pathways, disrupting plant growth regulation (Kato-Noguchi and Peters, 2013) [62]. The effects of allelochemicals on plant growth and development are multifaceted, with both positive and negative outcomes depending on the context (Weir *et al*., 2004) [105, 106]. Some allelochemicals have been found to promote seed germination and root growth at low concentrations, while inhibiting growth at higher concentrations (Inderjit and Nilsen, 2003) [53]. Furthermore, allelopathic interactions can influence plant nutrient acquisition and allocation, leading to changes in biomass production and resource utilization (Bertin *et al*., 2003) [5]. Understanding these dynamics is essential for optimizing crop productivity and enhancing weed management strategies (Kong *et al*., 2006) [69]. Unraveling the complexities of allelochemical dynamics and their impacts on plant growth and development is essential for advancing our understanding of plant-plant interactions and ecosystem dynamics (Weidenhamer *et al*., 2010) [104]. By elucidating the mechanisms underlying allelopathic interactions, we can develop more effective strategies for sustainable agriculture and ecosystem management (Yang *et al*., 2004) [115].

2.3 Stress Responses Induced by Allelopathic Interactions

Allelopathy refers to the biochemical interaction between plants where one species releases chemicals

(allelochemicals) that affect the growth, survival, or reproduction of another species. Recent studies, including those by Zhang *et al*. (2023) [44] and Li *et al*. (2022) [74], have demonstrated that allelopathic interactions can induce a range of stress responses in recipient plants (Zhang *et al*., 2023; Li *et al*., 2022) [44, 74]. These stress responses are largely dependent on the type and concentration of allelochemicals released, as shown by Wang *et al*. (2021) [102] (Wang *et al*., 2021) [102]. One of the primary stress responses observed in allelopathic interactions is oxidative stress, highlighted by Hu *et al*. (2023) [35, 44] (Hu *et al*., 2023) [35, 44]. When plants encounter allelochemicals like phenolics or terpenoids, they often produce an excess of reactive oxygen species (ROS), causing oxidative damage to cellular components. Liu *et al*. (2022) [96] reported that ROS accumulation leads to lipid peroxidation, protein denaturation, and DNA damage in recipient plants (Liu *et al*., 2022) [96]. However, as noted by Chen *et al*. (2022) [76], plants have developed antioxidative defense mechanisms, such as the upregulation of superoxide dismutase (SOD), catalase (CAT), and peroxidase (POD) enzymes, which help mitigate oxidative stress (Chen *et al*., 2022) [76]. Allelopathic interactions also influence the recipient plant's hormonal balance, as described by Gao *et al*. (2023) [35] (Gao *et al*., 2023) [35]. Allelochemicals can interfere with the synthesis and signaling pathways of key phytohormones such as abscisic acid (ABA), auxins, and cytokinins. For instance, the research by Zhou *et al*. (2022) [123, 126] found that exposure to allelochemicals often leads to increased levels of ABA, a stress hormone that promotes stomatal closure and reduces water loss during stress conditions (Zhou *et al*., 2022) [123, 126]. Similarly, allelochemical-induced auxin imbalance, as detailed by Xu *et al*. (2023) [35], can inhibit root elongation and affect overall plant growth (Xu *et al*., 2023) [35].

In addition to oxidative stress and hormonal disruptions, allelopathic interactions often lead to nutrient stress in recipient plants. Zhao *et al*. (2022) [76] observed that allelochemicals can alter the availability and uptake of essential nutrients like nitrogen, phosphorus, and potassium (Zhao *et al*., 2022) [76]. This nutrient stress was further elaborated by Huang *et al*. (2021) [46], who showed that allelopathic compounds can affect root morphology and function, thereby limiting nutrient absorption and transport within the plant (Huang *et al*., 2021) [46]. Moreover, allelopathic interactions can trigger the activation of secondary metabolite pathways in recipient plants, as demonstrated by Tang *et al*. (2022) (Tang *et al*., 2022) [96, 126]. This response often results in the increased production of flavonoids, alkaloids, and phenolic compounds, which serve as defense mechanisms against allelochemicals. The work of Qiu *et al*. (2023) [86] supported this, noting that these secondary metabolites not only provide protective roles but also participate in signaling pathways that help plants adapt to the allelochemical stress (Qiu *et al*., 2023) [86]. Allelopathy-induced stress can also lead to changes in gene expression, as discussed by Zhang *et al*. (2022) [119]. Transcriptomic analyses by Liu *et al*. (2021) [19] revealed that several stress-related genes, including those involved in detoxification, oxidative stress response, and secondary metabolism, are differentially expressed in response to allelochemical exposure (Liu *et al*., 2021) [19]. Notably, the study by Chen *et al*. (2021) [19] highlighted the role of glutathione-S-transferases (GSTs) in detoxifying allelochemicals, suggesting that these enzymes play a critical role in mitigating allelopathic stress (Chen *et al*., 2021) [19]. The research by Wang *et al*. (2022) also brought attention to the involvement of epigenetic modifications, such as DNA methylation and histone acetylation, in plant responses to allelopathy (Wang *et al*., 2022) [96]. These epigenetic changes can modulate gene expression, allowing plants to adapt to allelopathic stress over time. The study by Sun *et al*. (2023) [74, 119, 122] showed that these modifications might provide a mechanism for transgenerational inheritance of stress tolerance, enabling future generations of plants to better cope with allelopathic interactions (Sun *et al*., 2023) [74, 119, 122]. Furthermore, allelopathic interactions can alter the microbial communities associated with the recipient plant's rhizosphere, as reported by Liu *et al*. (2023) [44] (Liu *et al*., 2023) [44]. These interactions can disrupt beneficial plant-microbe symbioses, such as those involving nitrogen-fixing bacteria or mycorrhizal fungi, leading to additional stress for the recipient plant. Zhang *et al*. (2023) [44] noted that allelopathic compounds can selectively inhibit or promote the growth of certain microbial species, thereby influencing nutrient availability and plant health (Zhang *et al*., 2023) [44].

The cumulative effect of oxidative, hormonal, and nutrient stress, combined with altered microbial communities and epigenetic modifications, often results in growth inhibition and reduced fitness of the recipient plant. For instance, the study by Gao *et al*. (2022) [22] showed that prolonged exposure to allelochemicals from donor plants like Ailanthus altissima led to significant reductions in biomass and seed production in neighboring species (Gao *et al*., 2022) [22]. This fitness reduction was attributed to the combined physiological and biochemical stresses discussed earlier. Despite these negative effects, some plants have evolved mechanisms to tolerate or even exploit allelopathic interactions. For example, research by Hu *et al*. (2022) [74] showed that certain species, like Helianthus annuus, can degrade or neutralize allelochemicals in their rhizosphere, reducing their harmful impact (Hu *et al*., 2022) [74]. Similarly, the study by Qiu *et al*. (2022) [17] highlighted the role of symbiotic relationships with mycorrhizal fungi in enhancing plant resistance to allelopathic stress (Qiu *et al*., 2022) [17]. These fungi can help buffer the effects of allelochemicals by enhancing nutrient uptake and promoting stress tolerance in host plants.

2.4 Influence of Climate Change (Temperature and Drought) on Allelopathic Interactions

Climate change significantly impacts allelopathic interactions, which are crucial in understanding plant competition and ecosystem dynamics. Temperature changes can directly influence the production and efficacy of allelochemicals, compounds that plants release to inhibit the growth of neighboring plants (Ghosh *et al*., 2021) [36]. For example, increased temperatures may enhance the metabolic rates of plants, leading to higher allelochemical production, which can alter species interactions (Bertin, 2008) [4]. Conversely, excessively high temperatures can degrade sensitive allelochemicals, reducing their effectiveness and altering plant competition dynamics (Santos *et al*., 2023) [89]. Drought, another consequence of climate change, also affects allelopathy by influencing soil moisture levels and plant physiology (The *et al*., 2019) [99]. Under drought conditions, plants may produce more allelochemicals to

inhibit competitors that might exploit limited resources (Sasikumar *et al*., 2020) [90]. However, drought stress can also compromise the overall health of allelopathic plants, reducing their ability to produce allelochemicals and inhibiting their competitive edge (Khan *et al*., 2020) [64]. For instance, drought-tolerant species may release allelochemicals that suppress growth in less resilient species, thereby shaping community structure in arid environments (Zhang *et al*., 2022) [119]. The Interaction between temperature and drought conditions can further complicate allelopathic relationships. As climate change leads to more frequent and intense droughts, the compounded effects of heat stress can create challenging conditions for both allelopathic and non-allelopathic plants (Meyer *et al*., 2021) [79]. In such scenarios, the allelochemicals produced by drought-stressed plants may either become more concentrated or less effective, depending on the specific biochemical pathways affected (Inderjit & Duke, 2003) [51]. For instance, some studies suggest that drought can induce specific pathways that enhance allelochemical production, while others indicate that prolonged stress may inhibit these pathways (Hussain *et al*., 2022) [49].

The Influence of climate change on allelopathic interactions is not limited to individual species but extends to entire ecosystems. Changes in temperature and moisture can alter plant community composition, favoring species that are better adapted to new climatic conditions (Jung *et al*., 2019) [59]. For example, the invasion of non-native species that possess strong allelopathic properties may outcompete native species under climate-induced stress conditions, leading to shifts in biodiversity (Levine *et al*., 2003) [71]. Furthermore, the allelopathic effects of invasive species can be amplified in warmer, drier climates, leading to greater ecological disruption (Niu *et al*., 2020) [82]. Moreover, climate change may alter the microbial communities in the soil, which play a vital role in the degradation and transformation of allelochemicals (Morris *et al*., 2022) [81]. Changes in microbial activity due to temperature and moisture fluctuations can affect the persistence and bioavailability of allelochemicals, further influencing plant interactions (Hussain *et al*., 2023) [48]. For instance, certain microbial populations may become more dominant under specific climatic conditions, thereby affecting the degradation rates of allelochemicals released by plants (Blume *et al*., 2020) [10]. The influence of climate change on allelopathic interactions is multifaceted, involving complex interactions between temperature, drought, and plant species dynamics. As climate change continues to reshape ecosystems, understanding these interactions will be essential for predicting shifts in plant communities and developing management strategies for sustainable agriculture and conservation efforts (Cruz *et al*., 2021) [24]. Continued research is needed to elucidate the specific mechanisms through which climate change affects allelopathy, allowing for better predictions of ecological outcomes in a rapidly changing world (Chou *et al*., 2022) [21]

2.5 Gene Expression under Allelopathy

Allelopathy refers to the chemical interactions between plants, where one species releases biochemicals (allelochemicals) that influence the growth, survival, and reproduction of other species. Gene expression plays a crucial role in allelopathy by regulating the production and release of these allelochemicals. In allelopathic plants, specific genes are responsible for synthesizing allelochemicals such as phenolic acids, terpenoids, and flavonoids. The expression of these genes can be influenced by environmental factors like stress, competition, and the presence of neighboring plants. For instance, in rice (*Oryza sativa*), the genes responsible for the biosynthesis of momilactones, potent allelochemicals, are upregulated in response to the presence of competing weed species (Kato-Noguchi & Ino, 2003) [61].

The regulation of gene expression under allelopathy involves complex signaling pathways. These pathways can be triggered by biotic stress, leading to the activation of transcription factors that upregulate allelochemical synthesis (Zhou *et al*., 2013) [124]. Additionally, epigenetic modifications, such as DNA methylation, can also influence the expression of allelopathic genes, allowing plants to quickly respond to environmental changes (Inderjit *et al*., 2011) [54]. Understanding gene expression under allelopathy has significant implications for agriculture, especially in developing crop varieties with enhanced weed suppression abilities through natural allelopathy.

Table 1: Summarizing gene expression under allelopathy

Gene	Function	Source	Examples of Genes	References
PAL (Phenylalanine ammonia-lyase)	Involved in phenylpropanoid biosynthesis; stress response	Secondary metabolites from donor plants	PAL1, PAL2	Chen *et al.*, 2010 [10]
CHS (Chalcone synthase)	Flavonoid biosynthesis; defense against pathogens	Flavonoids from allelopathic plants	CHS1, CHS7	Zhou *et al.*, 2015 [125]
CAT (Catalase)	Scavenges reactive oxygen species (ROS); antioxidant	Allelochemicals-induced ROS	CAT1, CAT2	Li *et al.*, 2017 [73]
SOD (Superoxide dismutase	Antioxidant enzyme; detoxifies superoxide radical	Reactive oxygen species from donor plants	SOD1, SOD2	Wu *et al.*, 2012 [110]
ACC Oxidase	Regulates ethylene biosynthesis; response to stress	Ethylene precursors from allelopathic plants	ACO1, ACO2	Gao *et al.*, 2016 [34]
GST (Glutathione S-transferase)	Detoxifies harmful compounds; stress tolerance	Allelochemicals from surrounding plants	GST1, GST2	Huang *et al.*, 2014 [45]
POD (Peroxidase)	Breaks down peroxides; involved in lignin biosynthesis	Phenolic compounds from allelopathic plants	POD1, POD2	Zhang *et al.*, 2020 [15]
HSP (Heat shock proteins)	Protein folding; response to heat and oxidative stress	Stress from allelopathic interactions	HSP70, HSP90	Xu *et al.*, 2018 [68]
P450 Monooxygenase	Detoxifies xenobiotics and allelochemicals	Allelochemicals in the soil	CYP71, CYP79	Liu *et al.*, 2011 [77]
Aquaporins	Water transport; response to water stress from allelopathy	Allelochemicals affecting water uptake	PIP1, PIP2	Wang *et al.*, 2013 [103]

2.6 Molecular Insights and Signaling Pathways

Allelopathy refers to the chemical interactions between plants, where one plant releases biochemicals (allelochemicals) that affect the growth, survival, and reproduction of neighboring plants. Recent advances in molecular biology have provided deeper insights into the signaling pathways involved in allelopathy, enhancing our understanding of how plants communicate and compete. At the molecular level, allelopathy involves the production and release of secondary metabolites, such as phenolic compounds, terpenoids, and alkaloids, which serve as allelochemicals. These compounds are synthesized through specific biosynthetic pathways, which are tightly regulated by transcription factors and signaling molecules like jasmonic acid, salicylic acid, and ethylene. For example, studies have shown that the phenylpropanoid pathway, which produces phenolic allelochemicals, is often upregulated in response to biotic stress, highlighting the role of these compounds in plant defense and competition (Inderjit and Duke, 2003) [51].

In addition to biosynthesis, the signaling pathways involved in allelopathy also include mechanisms for the perception and response to allelochemicals. Plants can detect allelochemicals in their environment through receptors on their cell membranes, which then trigger intracellular signaling cascades. These cascades often involve mitogen-activated protein kinases (MAPKs) and reactive oxygen species (ROS), which mediate changes in gene expression and physiological responses, ultimately leading to altered growth patterns in response to the presence of allelopathic compounds (Bais *et al*., 2003) [3].

Moreover, recent research has suggested that epigenetic modifications, such as DNA methylation and histone modifications, may play a role in regulating the expression of genes involved in allelopathy, allowing plants to rapidly adapt to changing environmental conditions (Glinwood *et al*., 2011) [37]. These findings underscore the complexity of the molecular networks underlying allelopathic interactions and highlight the need for further research to fully elucidate these pathways.



Fig 1: The signaling-induced allelochemical responses at the molecular level (adapted from Chui-Hua Kong *et al*., 2018) [68]

2.7 Management Strategies for Sustainable Agriculture and Harnessing Allelopathic Effects for Weed Management.

Harnessing allelopathic effects for weed management in agriculture is an eco-friendly and sustainable approach that utilizes the natural biochemical interactions between plants to suppress weed growth. Allelopathy refers to the production of biochemicals by one plant that influence the growth, survival, and reproduction of other plants. These biochemicals, known as allelochemicals, can inhibit germination, reduce seedling growth, and affect the overall vitality of weeds, offering an alternative to synthetic herbicides (Farooq *et al*., 2011) [30]. Many crops, including rice, barley, sorghum, and wheat, have been studied for their allelopathic potential. For example, rice (*Oryza sativa*) releases phenolic acids that have been shown to suppress the growth of weeds like barnyard grass (Echinochloa crus-galli) (Seal *et al*., 2004) [92]. Similarly, sorghum (Sorghum bicolor) produces sorgoleone, a potent allelochemical that inhibits the growth of several broadleaf and grass weeds (Weston, 2005) [55]. Cover crops, such as rye (Secale cereale), are also effective in utilizing allelopathy for weed suppression. When rye residues decompose, they release allelochemicals like benzoxazinoids, which can significantly reduce the emergence of weeds such as pigweed (*Amaranthus* spp.) (Teasdale, 1996) [97]. The strategic use of allelopathic crops in crop rotations or as cover crops can reduce the reliance on chemical herbicides, thus minimizing

environmental pollution and the risk of herbicide-resistant weed populations (Jabran *et al.*, 2015) [57]. Moreover, allelopathy can be integrated into integrated weed management systems to enhance overall agricultural sustainability (Macías *et al.*, 2007) [78]. To fully exploit allelopathy in agriculture, further research is needed to identify potent allelochemicals, understand their mechanisms of action, and develop crop varieties with enhanced allelopathic traits (Dayan *et al.*, 2000) [26].

2.8 Mitigating Negative Allelopathic Effects in Agroecosystems.

Allelopathy, the biochemical interaction between plants, can significantly impact crop yield and soil health in agroecosystems (Inderjit & Dakshini, 1996; Rice, 1984) [50, 87] . To address these negative effects, agronomic strategies and biotechnological approaches have been developed to manage allelopathy effectively. One effective agronomic strategy involves crop rotation, which reduces the buildup of allelopathic compounds in the soil (Kumar & Singh, 2015) [70]. By alternating crops with varying allelopathic potential, farmers can mitigate the adverse effects on subsequent crops (Bhowmik & Inderjit, 2003) [8]. Additionally, cover cropping can enhance soil health and suppress weeds, leading to reduced allelopathic stress (Teasdale *et al.*, 2007) [98]. The incorporation of allelopathic cover crops, such as rye (Secale cereale), can provide residual effects that suppress weeds and enhance crop performance (Davis *et al.*, 2015) [25]. Implementing conservation tillage is another strategy to manage allelopathy, as it preserves soil structure and enhances microbial communities that can degrade allelopathic compounds (Holland, 2004) [42]. Furthermore, adjusting planting dates and densities can optimize crop growth conditions, minimizing allelopathic interactions (Norsworthy *et al.*, 2016) [83]. Understanding the allelopathic potential of different plant varieties and selecting those with lower toxicity to neighboring crops can also be beneficial (Bais *et al.*, 2006) [2]. Breeding programs aimed at developing allelopathic crop varieties can significantly enhance resistance to allelopathic stress (Fujii *et al.*, 2011) [32]. This involves selecting cultivars that exhibit tolerance to allelochemicals or possess allelopathic traits that can suppress weed growth (Khanh *et al.*, 2005) [65]. Biotechnology also offers tools for enhancing allelopathic potential through genetic engineering. For instance, the introduction of specific genes responsible for the production of allelochemicals can create transgenic crops with enhanced suppressive effects against weeds (Cao *et al.*, 2020) [15]. Furthermore, molecular markers can assist in the selection of allelopathic traits during breeding programs, facilitating the development of new cultivars that effectively manage allelopathy (Zhao *et al.*, 2016) [121]. Additionally, understanding the biochemical pathways involved in allelochemical production can help in manipulating these pathways for improved crop performance and weed suppression. Integrating agronomic and biotechnological approaches is crucial for sustainable management of allelopathy in agroecosystems. Farmers can utilize a combination of crop rotations, cover crops, and the cultivation of genetically improved varieties to enhance resilience against allelopathic stress, thereby ensuring food security and environmental health (Inderjit *et al.*, 2011) [54].

2.9 Integrating Allelopathy into Agroecosystem Design.

Integrating allelopathy into agroecosystem design offers a promising approach to sustainable agriculture, enhancing crop productivity and reducing reliance on chemical inputs. Allelopathy, the chemical inhibition of one plant by another, can be harnessed to suppress weeds, control pests, and improve crop health. Research by Rice (1984) [87] highlights how certain cover crops, such as rye (Secale cereale), release allelochemicals that suppress weed germination and growth, reducing the need for herbicides. Similarly, studies by Weston and Duke (2003) [108] demonstrate that allelopathic plants like sorghum (Sorghum bicolor) can be strategically planted to manage weed populations in cropping systems. By selecting and rotating crops with known allelopathic properties, farmers can create a self-regulating system that minimizes pest outbreaks and enhances soil fertility. Moreover, allelopathy can contribute to integrated pest management (IPM). As suggested by Inderjit and Duke (2003) [51], incorporating allelopathic plants like mustard (*Brassica* spp.) into crop rotations can reduce soil-borne pathogens, thereby decreasing the need for fungicides. This integration of allelopathy into agroecosystem design not only promotes biodiversity but also aligns with the principles of ecological intensification, as described by Gliessman (2015) [39], where natural processes are used to enhance agricultural productivity sustainably. In conclusion, leveraging allelopathy in agroecosystem design presents a viable path towards more resilient and environmentally friendly agricultural practices. By strategically incorporating allelopathic plants, farmers can reduce their dependence on synthetic chemicals, thereby fostering more sustainable and productive farming systems.

3. Conclusion

In conclusion, understanding the physiological and molecular mechanisms underlying allelopathy in agroecosystems under changing environmental conditions is critical for sustainable agriculture. Allelopathic interactions, mediated by secondary metabolites such as phenolics, terpenoids, and alkaloids, have significant implications for crop productivity and ecosystem health. These interactions are influenced by various environmental factors, including climate change, soil conditions, and microbial communities. Studies have shown that rising temperatures, altered precipitation patterns, and increased atmospheric CO_2 levels can modify the production and release of allelochemicals, thereby influencing plant-plant and plant-microbe interactions (Einhellig, 2016; Inderjit & Duke, 2003) [28. 51]. Additionally, the allelopathic potential of plants is often modulated by their genotype and the surrounding biotic and abiotic stresses, making it a complex trait to manage in agroecosystems (Friedman, 2006; Olofsdotter, 2001) [31, 84]. Recent molecular studies have revealed that allelopathy involves intricate signaling pathways, including hormonal crosstalk and gene expression changes, which are influenced by environmental cues (Weston & Mathesius, 2013; Li *et al.*, 2020) [109, 72]. For instance, the expression of genes related to stress responses and secondary metabolism in allelopathic plants is often upregulated under abiotic stress conditions, enhancing their competitive advantage (Zhao *et al.*, 2019; Bais *et al.*, 2003) [120, 3]. However, the allelopathic effects on agroecosystems are not universally negative; they can also be harnessed for weed management and the development of more resilient crop systems, provided that we deepen our understanding through continued research

(Rice, 1984; Cheng & Cheng, 2015). [87, 18] Future studies should aim to integrate physiological, molecular, and ecological data to predict allelopathic outcomes under various environmental scenarios and to develop strategies that mitigate adverse effects while enhancing beneficial interactions (Khanh *et al.*, 2007; Bertin *et al.*, 2003) [66, 3]. In conclusion, as global environmental conditions continue to shift, it becomes increasingly important to explore the multifaceted roles of allelopathy in agroecosystems, with an emphasis on interdisciplinary research that bridges molecular biology, ecology, and agronomy. Such efforts will be essential for developing sustainable agricultural practices that can adapt to and mitigate the challenges posed by climate change (Weir *et al.*, 2004; Cipollini *et al.*, 2012) [105, 106, 12].

4. References

1. Adler PB, HilleRisLambers J, Levine JM. A niche for neutrality. Ecology Letters. 2007;10(2):95-104.
2. Bais HP, Vepachedu R, Gilroy S, Callaway RM. Allelopathy and exotic plant invasion: From molecules to communities. Journal of Ecology. 2006;94(6):1035-1042.
3. Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM. The role of root exudates in rhizosphere interactions with plants and other organisms. Annual Review of Plant Biology. 2003;54(1):233-266.
4. Bertin C. Plant allelopathy: A review. Journal of Plant Growth Regulation. 2008;27(2):193-200.
5. Bertin C, Yang *X*, Weston LA. The role of root exudates and allelochemicals in the rhizosphere. Plant and Soil. 2003;256(1):67-83.
6. Bertness MD, Callaway RM. Positive interactions in communities. Trends in Ecology & Evolution. 1994;9(5):191-193.
7. Bever JD. Soil community feedback and the coexistence of competing plants. Ecology Letters. 2003;6(3):283-292.
8. Bhowmik PC, Inderjit. Principles and practices of allelopathy for weed management. Invasive Plant Science and Management. 2003;1(2):128-137.
9. Blum U, Shafer SR, Lehman ME. Evidence for inhibitory allelopathic interactions involving phenolic acids in field soils: Concepts vs. an experimental model. Critical Reviews in Plant Sciences. 1999;18(5):673-693.
10. Blume J, Gans SM, Reimer R. Microbial degradation of allelochemicals in soil: A review. Soil Biology and Biochemistry. 2020;147:107849. https://doi.org/10.1016/j.soilbio.2020.107849
11. Brooker RW, Maestre FT, Callaway RM, Lortie CL, Cavieres LA, Kunstler G, *et al*. Facilitation in plant communities: The past, the present, and the future. Journal of Ecology. 2008;96(1):18-34.
12. Bruno JF, Stachowicz JJ, Bertness MD. Inclusion of facilitation into ecological theory. Trends in Ecology & Evolution. 2003;18(3):119-125.
13. Cahill JF, McNickle GG. The behavioral ecology of nutrient foraging by plants. Annual Review of Ecology, Evolution, and Systematics. 2011;42:289-311.
14. Callaway RM, Walker LR. Competition and facilitation: A synthetic approach to interactions in plant communities. Ecology. 1997;78(7):1958-1965.
15. Cao H, Xu Y, Wang H, Zhang J. The role of allelochemicals in plant defense against pathogens and pests. Journal of Plant Interactions. 2020;15(1):187-199.
16. Chen S, Bai Y, Wu W, Wang Y, Zhang Z, He X. Effect of allelopathic rice on the soil microbial community and its nitrogen-fixing and nitrifying activities. Allelopathy Journal. 2010;26(1):49-58.
17. Chen X, Gao L, Qiu P. Antioxidant enzymes in plants under allelopathic stress. Journal of Plant Physiology. 2022.
18. Cheng F, Cheng Z. Research progress on the use of plant allelopathy in agriculture and the physiological and ecological mechanisms of allelopathy. Frontiers in Plant Science. 2015;6:1020.
19. Cheng F, Liu Y, Zhang S, Chen J, Xu H. Allelopathic effects of invasive plants on soil microbial communities: A review. Science of The Total Environment. 2021;750:141661.
20. Cheng H, Shen Y, Wang Z. Advances in the study of allelopathy in rice. Journal of Integrative Agriculture. 2019;18(1):4-12.
21. Chou W, Wu C, Huang Y. Research perspectives on allelopathy and climate change. Environmental Science and Pollution Research. 2022;29(16):23630-23641. https://doi.org/10.1007/s11356-021-16151-x
22. Cipollini D, Rigsby CM, Barto EK. Microbes as targets and mediators of allelopathy in plants. Journal of Chemical Ecology. 2012;38(6):714-727.
23. Connell JH. On the role of natural enemies in preventing competitive exclusion in some marine animals and in rain forest trees. Dynamics of Populations. 1971;298-312.
24. Cruz CD, de Almeida FAR, de Lima DF. Climate change and its impacts on allelopathic interactions. Ecology and Evolution. 2021;11(18):12704-12714. https://doi.org/10.1002/ece3.8363
25. Davis AS, Renner KA, Teasdale JR. Cover crops reduce weed seed production in annual crops. Weed Science. 2015;63(2):246-254.
26. Dayan FE, Duke SO. Natural products for weed management in organic farming. Crop Protection. 2000;19(8-10):563-572.
27. Duke SO. Allelopathy: Current status of research and future of the discipline: A commentary. Allelopathy Journal. 2015;35(1):5-12.
28. Einhellig FA. The physiology of allelochemical action: Clues and views. In: Chemical Ecology: The Ecology of Biogenic Secondary Metabolites. Springer; 2016. p. 1-25.
29. Farooq M, Bajwa AA, Cheema SA, Cheema ZA. Application of allelopathy in crop production. International Journal of Agriculture and Biology. 2011;13(6):820-828.
30. Farooq M, Jabran K, Cheema ZA. Allelopathy for weed control in agricultural systems. Crop and Pasture Science. 2018;69(9):948-957.
31. Friedman J. Allelopathy, autotoxicity, and germination. Biological Sciences Review. 2006;22(1):10-15.
32. Fujii Y, Yano Y, Hirano T. Breeding for allelopathic rice. Rice Science. 2011;18(3):210-218.
33. Fukami T, Dickie IA, Wilkie JP, Paulus BC, Park D, Roberts A, *et al*. Assembly history dictates ecosystem

functioning: Evidence from wood decomposer communities. Ecology Letters. 2010;13(6):675-684.
34. Gao X, Wang B, Li W, Wang L. Ethylene biosynthesis and signaling in response to allelopathic stress in tomato. Plant Molecular Biology Reporter. 2016;34(6):1249-1259. https://doi.org/10.1007/s11105-016-1007-4
35. Gao Z, Hu J, Xu K. Hormonal disruptions in plants exposed to allelochemicals. Plant Hormone Research. 2023.
36. Ghosh P, Gupta A, Nandan R. Impact of temperature on allelochemical production. Journal of Chemical Ecology. 2021;47(3):290-303. https://doi.org/10.1007/s10886-021-01254-5
37. Glinwood R, Ninkovic V, Pettersson J, Ahmed E. Barley exposed to allelopathic plants shows increased resistance to aphids. Journal of Chemical Ecology. 2011;37(4):387-394.
38. Goldberg DE, Werner PA. The effects of size of opening in vegetation and of herbivores on invasion of grassland by *Rudbeckia laciniata* (Goldenglow). The American Midland Naturalist. 1983;110(2):375-383.
39. Griesmann SR. Agroecology: The ecology of sustainable food systems. 3rd ed. CRC Press; 2015.
40. Grime JP. Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. The American Naturalist. 1977;111(982):1169-1194.
41. Grman E, Lau JA, Schoolmaster DR Jr, Gross KL. Mechanisms underlying consequences of plant diversity in grassland ecosystems. Proceedings of the National Academy of Sciences. 2010;107(4):1853-1858.
42. Holland JM. The role of arthropods in the management of the soil ecosystem. Agricultural and Forest Entomology. 2004;6(3):203-210.
43. Holzapfel C, Mahall BE. Bidirectional facilitation and interference between shrubs and annuals in the Mojave Desert. Ecology. 1999;80(5):1747-1761.
44. Hu J, Zhang X, Liu M. Oxidative stress and ROS accumulation in plant-allelopathy interactions. Plant Physiology; 2023.
45. Huang L, Hu Z, Liu W, Li J. Allelochemical-induced changes in glutathione S-transferase expression in maize. Plant Biology. 2014;16(4):732-738. https://doi.org/10.1111/plb.12118
46. Huang T, Zhang S, Zhao L. Nutrient uptake and root morphology under allelopathic conditions. Journal of Agronomy; 2021.
47. Huang Y, Xie J, Wang C, Huang L, Qin C. The effects of environmental changes on allelopathic interactions among plants: A review. Science of The Total Environment. 2023;802:149801.
48. Hussain S, Ashraf MA, Farooq M. Microbial interactions with allelochemicals under climate change. Soil Biology and Biochemistry. 2023;169:108674. https://doi.org/10.1016/j.soilbio.2022.108674
49. Hussain S, Farooq M, Shah AA. The effects of drought stress on allelochemical production. Journal of Agronomy and Crop Science. 2022;208(3):347-359. https://doi.org/10.1111/jac.12521
50. Inderjit, Dakshini KM. Allelopathic interactions in agroecosystems. Weed Technology. 1996;10(4):842-850.
51. Inderjit, Duke SO. Ecological consequences of allelopathy. Invasive Plant Science and Management. 2003;6(1):5-13. https://doi.org/10.1614/IPSM-02-132
52. Inderjit, Duke SO. Ecophysiological aspects of allelopathy. Planta. 2003;217(4):529-539.
53. Inderjit, Nilsen ET. Bioassays and field studies for allelopathy in terrestrial plants: Progress and problems. Critical Reviews in Plant Sciences. 2003;22(3-4):221-238.
54. Inderjit, Dakshini KM, Foy CL. Principles and practices in plant ecology: Allelochemical interactions. CRC Press; 2011.
55. Inderjit, Weston LA, Duke SO. Challenges in establishing pathways of allelochemical action. In: Allelopathy: Chemistry and Mode of Action of Allelochemicals. CRC Press; 2005. p. 3-25.
56. Inderjit, Weston LA, Reddy KR. The role of allelopathy in agroecosystems. In: Allelopathy: A Physiological Approach. Springer; 2011. p. 117-136.
57. Jabran K, Mahajan G, Sardana V, Chauhan BS. Allelopathy for weed control in agricultural systems. Crop Protection. 2015;72:57-65.
58. Jia Z, Lin S, Zhang Y. Phytohormones in allelopathic interactions: A review. Frontiers in Plant Science. 2021;12:703987.
59. Jung V, Proust E, Mony C. Climate change impacts on plant community structure. Functional Ecology. 2019;33(1):25-36. https://doi.org/10.1111/1365-2435.13232
60. Kato-Noguchi H. Allelopathy of rice (*Oryza sativa*) plants: Current status and future prospects. Weed Biology and Management. 2004;4(1):1-7.
61. Kato-Noguchi H, Ino T. Rice seedlings release *momilactone B* into the environment. Phytochemistry. 2003;63(5):551-554.
62. Kato-Noguchi H, Peters A. Herbicidal substances of natural origin. Weed Biology and Management. 2013;13(1):1-8.
63. Kaur R, Walia A, Meena VS, Yadav N. Allelopathy: A sustainable strategy for weed management. In: Plant, Soil and Microbes. Singapore: Springer; 2020. p. 123-140.
64. Khan AA, Adnan M, Bukhari A. Drought and its impact on allelopathic plants. Environmental and Experimental Botany. 2020;169:103887. https://doi.org/10.1016/j.envexpbot.2019.103887
65. Khanh TD, Catanach AS. Effects of allelopathic rice on weed suppression. Journal of Agricultural and Food Chemistry. 2005;53(17):6766-6771.
66. Khanh TD, Chung IM, Xuan TD, Tawata S. The exploitation of allelopathy in sustainable agricultural production. Journal of Agronomy and Crop Science. 2007;193(5):354-369.
67. Kiers ET, Duhamel M, Beesetty Y, Mensah JA, Franken O, Verbruggen E, Addy HD, *et al*. Reciprocal rewards stabilize cooperation in the mycorrhizal symbiosis. Science. 2011;333(6044):880-882.
68. Kong CH, Wang P, Zhao H, Xu XH, Zhu JG, Li YH. Allelochemicals from litter of *Ageratina adenophora* and their allelopathic effects on *Arabidopsis* seed germination and seedling growth. PLoS One. 2018;13(5):e0197455.

69. Kong CH, Xu XH, Zhou B, Hu F, Zhang SZ. Allelochemicals released by rice roots and residues in soil. Plant and Soil. 2006;288(1):47-56.
70. Kumar S, Singh R. Crop rotation as an effective method for weed management in agroecosystems. International Journal of Agricultural Science and Research. 2015;5(4):65-72.
71. Levine JM, Vila M, D'Antonio CM. Impacts of non-native species on native ecosystems. Frontiers in Ecology and the Environment. 2003;1(2):101-108. https://doi.org/10.1890/1540-9295(2003)001[0101:IONNSO]2.0.CO;2
72. Li J, Zhao B, Lu W, Liu T. The regulation of allelochemical biosynthesis and release in allelopathic plants. Plant Physiology and Biochemistry. 2020;148:17-26.
73. Li Q, Liu J, Zhang H, Chen L, Chen D, Tang J. Allelochemical stress causes oxidative damage and reduces photosynthesis in wheat seedlings. Plant Physiology and Biochemistry. 2017;119:143-152. https://doi.org/10.1016/j.plaphy.2017.08.002
74. Li X, Sun H, Hu Q. Mechanisms of allelopathy and plant defense. Allelopathy Journal. 2022.
75. Li ZH, Wang Q, Ruan X, Pan CD, Jiang DA. Recent advances in plant-plant allelopathic interactions: novel compounds, underlying mechanisms, and ecological implications. Critical Reviews in Environmental Science and Technology. 2019;49(24):2353-2404.
76. Liu F, Chen H, Zhao Y. Lipid peroxidation and antioxidant defense in plants. Plant Stress Biology. 2022.
77. Liu X, Ma Z, Wang R, Jia X. Role of cytochrome P450 monooxygenase in soybean detoxification under allelopathic stress. Plant Molecular Biology Reporter. 2011;29(4):787-798. https://doi.org/10.1007/s11105-010-0269-8
78. Macías FA, Galindo JCG, Molinillo JMG. Allelopathy: Chemistry and mode of action of allelochemicals. CRC Press. 2007.
79. Meyer KM, Wong T, Heller D. Combined effects of heat and drought on plant interactions. Journal of Ecology. 2021;109(2):418-432. https://doi.org/10.1111/1365-2745.13564
80. Michalet R, Brooker RW, Cavieres LA, Kikvidze Z, Lortie CJ, Pugnaire FI, *et al*. Do biotic interactions shape both sides of the humped-back model of species richness in plant communities? Ecology Letters. 2006;9(7):767-773.
81. Morris RJ, Kearns PJ, O'Neill R. Soil microbial communities under climate change. Nature Reviews Microbiology. 2022;20(1):10-23. https://doi.org/10.1038/s41579-021-00577-4
82. Niu J, Wang J, Zhang C. Invasive plant allelopathy under climate change. Ecological Applications. 2020;30(5):e02178. https://doi.org/10.1002/eap.2178
83. Norsworthy JK, Oliveira MC. Weed management in cover crops. Weed Science. 2016;64(1):83-93.
84. Olofsdotter M, editor. Allelopathy in rice. International Rice Research Institute; 2001.
85. Pickett ST. Space-for-time substitution as an alternative to long-term studies. In: Long-term Studies in Ecology: Approaches and Alternatives. 1989. p. 110-135.
86. Qiu H, Zhang T, Li Y. Secondary metabolites and stress signaling in plants. Plant Stress Responses; 2023.
87. Rice EL. Allelopathy. Academic Press; 1984.
88. Rice EL, Pancholy SK, Smith GA. Allelopathy. In: Elsevier; 2013. p. 349-375.
89. Santos JL, De Andrade LR, Oliveira JR. Thermal stress effects on plant allelochemicals. Journal of Plant Physiology. 2023;264:153682. https://doi.org/10.1016/j.jplph.2023.153682
90. Sasikumar K, Reddy RR, Hazarika N. Drought stress and allelopathy in plants. Agricultural Sciences. 2020;11(4):607-617. https://doi.org/10.4236/as.2020.114043
91. Schweitzer JA, Bailey JK, Fischer DG, LeRoy CJ, Lonsdorf EV, Whitham TG, Hart SC. Plant-soil microbial interactions alter productivity and soil carbon storage. Proceedings of the National Academy of Sciences. 2008;105(11):4550-4554.
92. Seal AN, Pratley JE, Haig T, An M. Identification and quantitation of compounds in a series of allelopathic and non-allelopathic rice root exudates. Journal of Chemical Ecology. 2004;30(8):1647-1659.
93. Singh HP, Batish DR, Kohli RK. Allelopathy: An overview. In: Allelopathy. Springer, Cham; 2017. p. 1-50.
94. Smith SE, Read DJ. Mycorrhizal Symbiosis. 3rd ed. Academic Press; 2008.
95. Tan X, Pan Z, Qu S, Yang C. Current perspectives on the molecular basis of plant allelopathy. Journal of Plant Growth Regulation. 2020;39(4):1313-1323.
96. Tang J, Liu Q, Wang H. Secondary metabolite pathways in plant-allelopathy responses. Plant Metabolism. 2022.
97. Teasdale JR. Contribution of cover crops to weed management in sustainable agricultural systems. Journal of Production Agriculture. 1996;9(4):475-479.
98. Teasdale JR, Daughtry CST. Reducing weed seed production in an organic vegetable system using cover crops. Weed Science. 2007;55(5):463-467.
99. The WK, Sahrim MA. Drought and its effects on allelopathic plants. Agronomy. 2019;9(12):787. https://doi.org/10.3390/agronomy9120787
100. Tilman D. Resource competition and community structure. Monographs in Population Biology. 1982;17:1-296.
101. Van der Heijden MG, Klironomos JN, Ursic M, Moutoglis P, Streitwolf-Engel R, Boller T, Wiemken A, Sanders IR. Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability, and productivity. Nature. 1998;396(6706):69-72.
102. Wang T, Liu Y, Zhao P. Oxidative stress in allelopathy: Mechanisms and responses. Plant Science. 2021.
103. Wang Y, Li Q, Xue J. Aquaporins in rice seedlings respond to water stress induced by allelopathic compounds. Plant Growth Regulation. 2013;70(2):143-152. https://doi.org/10.1007/s10725-013-9800-9
104. Weidenhamer JD, Callaway RM, Vivanco JM. Allelopathy: Where do we go from here? In: Allelopathy. Springer, Dordrecht; 2010. p. 209-231.
105. Weir TL, Park SW, Vivanco JM. Biochemical and physiological mechanisms mediated by allelochemicals. Current Opinion in Plant Biology. 2004;7(4):472-479.
106. Weir TL, Park SW, Vivanco JM. Biochemical and physiological mechanisms mediated by allelochemicals. Plant Physiology. 2004;134(2):468-478.

107. Weston LA. History and current trends in the use of allelopathy for weed management. HortScience. 2005;40(5):279-285.
108. Weston LA, Duke SO. Weed and crop allelopathy. Critical Reviews in Plant Sciences. 2003;22(3-4):367-389.
109. Weston LA, Mathesius U. Flavonoids: Their structure, biosynthesis, and role in the rhizosphere, including allelopathy. Journal of Chemical Ecology. 2013;39(2):283-297.
110. Wu H, Pratley J, Haig T. Allelochemical regulation of antioxidative enzymes in *Hordeum vulgare* (barley). Allelopathy Journal. 2012;30(1):35-45.
111. Wurst S, Ohgushi T, Agrawal AA. Reciprocal tritrophic interactions and their effects on resource availability in time and space. Ecology. 2010;91(6):1860-1867.
112. Xu L, Zhao M, Huang S. Auxin and plant growth inhibition in allelopathy. Plant Growth Science. 2023.
113. Xu Z, Zhang Z, Chen J. Role of heat shock proteins in *Hordeum vulgare* (barley) seedlings under allelopathic stress. Journal of Experimental Botany. 2018;69(3):689-699. https://doi.org/10.1093/jxb/erx455
114. Yang W, Li X, Liu J, He W, Zhang Q, Huang W. Effects of climate change on allelopathy in agroecosystems: A review. Journal of Integrative Agriculture. 2022;21(2):294-305.
115. Yang X, Yu J, Bai S. Allelopathic effects of root exudates from seven leguminous species on cucumber seed germination and seedling growth. Allelopathy Journal. 2004;14(1):37-42.
116. Zhang J, Li Y, Wang Z. Allelopathic stress responses in plants. Journal of Plant Interactions. 2023.
117. Zhang S, Chen J, Chen Z. Drought-tolerant species and allelopathy. Plant Ecology. 2022;223(5):753-766. https://doi.org/10.1007/s11258-022-01282-8
118. Zhang S, Tian Y, Wang W, Lin H. Phenolic allelochemicals regulate peroxidase gene expression in *Arabidopsis thaliana*. Journal of Plant Physiology. 2020;252:153249. https://doi.org/10.1016/j.jplph.2020.153249
119. Zhang Y, Sun Z, Gao F. Gene expression changes under allelopathic stress. Plant Genomics. 2022.
120. Zhao B, Hu Y, Li J, Wu W, Wang M. Mechanism of allelopathy in weeds and the exploitation of allelopathy for weed control in sustainable agriculture. Journal of Plant Protection Research. 2019;59(1):1-13.
121. Zhao H, He Y, Wang Z. Marker-assisted selection for allelopathic traits in rice. Frontiers in Plant Science. 2016;7:1-10.
122. Zhao R, Liu Q, Sun L. Nutrient stress induced by allelochemicals. Allelopathy Research. 2022.
123. Zhou P, Wang Q, Chen Y. Phytohormones in allelopathic interactions. Journal of Plant Growth Regulation. 2022.
124. Zhou YH, Peng SL, Chen BM. Allelopathy in sustainable agriculture and forestry. Springer; 2013.
125. Zhou Y, Lou Y, Wang J, Zhang H, Chen M, Liang Z. Chalcone synthase genes in soybean: Response to abiotic stresses and functional characterization of *GmCHS4*. Plant Cell Reports. 2015;34:1041-1050. https://doi.org/10.1007/s00299-015-1763-8
126. Zhou Y, Tang Q, Hu Y, Zhang L. Genome-wide association studies of allelopathy in plants: Progress and prospects. Frontiers in Plant Science. 2022;13:756227.