




International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
NAAS Rating (2025): 5.29
IJABR 2025; SP-9(10): 819-824
www.biochemjournal.com
Received: 10-08-2025
Accepted: 13-09-2025

NA Karishma
Department of Fruit Science,
College of Agriculture, Vellayani,
Thiruvananthapuram, Kerala,
India

S Simi
Department of Fruit Science,
College of Agriculture, Vellayani,
Thiruvananthapuram, Kerala,
India

PR Manju
Assistant Professor, Department
of Fruit Science, College of
Agriculture, Vellayani,
Thiruvananthapuram, Kerala,
India

B Rani
Professor and Head, Department
of Soil Science and Agricultural
Chemistry, College of
Agriculture, Vellayani
Thiruvananthapuram, Kerala,
India

Corresponding Author:
S Simi
Department of Fruit Science,
College of Agriculture, Vellayani,
Thiruvananthapuram, Kerala,
India

Growth response of guava (*Psidium guajava* L.) to differential spacing and nutrient management under high density planting system in Kerala

NA Karishma, S Simi, PR Manju, and B Rani

DOI: <https://www.doi.org/10.33545/26174693.2025.v9.i10Sj.5925>

Abstract

A study was conducted in guava to evaluate the effect of spacing and nutrient level and schedule on the vegetative parameters of guava under HDP system in Kerala. Experiment was laid out in split plot design with four levels of spacing as main plot and six fertilizer level as subplot treatments replicated thrice. The results indicated that plant height was significantly higher in closer spacing of 2.0 m × 1.5 m with 100% of fertilizer dose as per schedule-2 (S₄F₆; 1.23 m; 1.60 m) at 6 and 9 MAP. The highest plant girth (9.30 cm) and canopy spread in N-S direction (105.98 cm) were observed as a result of the interactive effect of 3.0 m × 3.0 m spacing with 100% of fertilizer dose as per schedule-2 (S₁F₆) at 6 MAP while at 9 MAP the highest values were obtained with 3.0 m × 3.0 m with 100% of fertilizer dose as per schedule-1 (S₁F₅) (13.15; 123.44 cm). Canopy spread in E-W direction was the highest in 3.0 m × 3.0 m spacing with 100% of fertilizer dose as per schedule-1 at 6 MAP (99.33 cm) and at 9 MAP 3.0 m × 3.0 m with 100% of fertilizer dose as per schedule-2 recorded the highest values (136.01 cm). Data on primary branch girth, secondary branch girth and pruned biomass showed that interactive effect has no significant effect at 6 and 9 MAP. The least number of days to shoot emergence (5.04) was observed in 3.0 m × 3.0 m with 100% of fertilizer dose as per schedule-2 at 9MAP. In conclusion, the growth parameters were observed to be better at highest spacing and nutrient dose (S₁F₆ and S₁F₅) except the plant height which was observed to be the highest with S₄F₆.

Keywords: Guava, Arka Kiran, plant height, spacing, canopy spread, plant girth, fertilizer, HDP

Introduction

Guava (*Psidium guajava* L.), popularly known as the “apple of the tropics,” ranks fourth in area and production after mango, banana, and citrus. Traditionally, guava is cultivated under conventional planting systems, which often hinder the attainment of optimal productivity. This limitation is primarily due to the large canopy size of the trees, resulting in lower yield per unit area and increased labour requirements (Singh *et al.*, 2003) ^[20]. In addition, such trees require several years to reach the bearing stage, thereby increasing the overall production cost per unit area. Maximizing productivity necessitates the adoption of modern and efficient production practices. Key strategies include utilizing high-quality planting material, optimizing planting density, implementing scientific canopy management techniques, and enhancing support and input management systems. High density planting not only enhances yield and economic returns per unit area during the initial years but also ensures more efficient utilization of available inputs (Reddy, 2004) ^[18]. Effective nutrient management plays a crucial role in sustaining productivity under high density systems. The nutrient requirement of guava varies across regions depending on soil characteristics and climatic conditions, which influence plant growth and flushing patterns. Guava is highly responsive to inorganic fertilizer application, and a balanced supply of nitrogen, phosphorus, and potassium is essential for optimal growth and fruit production (Kumar *et al.*, 2009) ^[11]. However, standardized recommendations for major nutrient application under high density planting system are yet to be established for local agroclimatic regions. Hence, the primary objective of the study was to evaluate the effect of spacing and nutrient levels and schedules on the vegetative parameters of guava under high density planting system in Kerala.

Materials and Methods

The field experiment was conducted at the Department of Fruit Science, College of Agriculture, Vellayani during the year 2023-2025. The primary objective of the study was to evaluate the effect of spacing and nutrient levels and schedules on vegetative parameters of guava under high density planting system. The area is situated at 8° 43' North Latitude and 76° 54' East Longitude at an altitude of 29 m above mean sea level. Four months old layers of guava variety Arka Kiran, a pink fleshed hybrid (Kamsari × Purple Local) released by the ICAR-Indian Institute of Horticultural Research (IIHR), Bengaluru was used for the study. Experiment was laid out in split plot design with four main plot treatments and six subplot treatments replicated three times. Main plot treatments included four levels of spacing (S); S₁-3.0 m × 3.0 m, S₂-3.0 m × 1.5 m, S₃-2.0 m × 2.0 m and S₄-2.0 m × 1.5 m. Subplot treatments included six levels of fertilizer dose and schedule (F); F₁-50% of fertilizer dose as per schedule-1, F₂-50% of fertilizer dose as per schedule-2, F₃-75% of fertilizer dose as per schedule-1, F₄-75% of fertilizer dose as per schedule-2, F₅-100% of fertilizer dose as per schedule-1 and F₆-100% of fertilizer dose as per schedule-2. Fertilizer doses were given based on KAU package of practices recommendations: Crops (KAU, 2016). For first year, 1/5th of RDF was given based on soil test data. In schedule 1 fertilizers were applied after each pruning in three equal splits. Table 1 indicates schedule 2.

Each subplot contained six plants which were maintained under uniform condition of orchard management system during the study period. All agronomic practices were carried out as per package of practices of KAU. First pruning was done at around 3 MAP maintaining a height of 45 to 60 cm, 3-4 primary branches were retained, second pruning was done after three months to primary branches by removing 50% of their length. Three secondary branches were retained. Third pruning was given after 3 months to secondary branches by removing 50% of their length. Three tertiary branches were retained.

Observations recorded

Vegetative parameters such as plant height (m) was measured using measuring tape in centimeters from soil level to the tip of the growing point and average was calculated, stem girth (cm) of the plant was recorded as circumference in centimeters 5-10 cm above soil level using a measuring tape and average was calculated, canopy spread (North-South direction (cm), East-West direction (cm)) was determined by measuring the widest horizontal distance between the outermost branches of each plant, encompassing the entire canopy using a measuring scale, circumference of the primary and secondary branches were measured and expressed in centimeters (cm), and pruned biomass (kg) was measured using weighing balance after each pruning and days to shoot emergence after each pruning was counted.

Results and Discussion

Plant height: The tallest plants were observed under S₁-3.0 m × 3.0 m (0.66 m) at 3 MAP and it was on par with S₄-2.0 m × 1.5 m (0.65 m), S₄ spacing at 6 MAP (1.08 m) and 9 MAP (1.42 m). The least plant height was observed in S₁ (0.83m and 1.17m respectively) at 6 MAP and 9 MAP. Among different fertilizer doses, the tallest plants were observed in F₂ (0.61 m) at 3 MAP. While at 6 MAP and 9

MAP the highest plant height was observed in F₆-100% of fertilizer dose as per schedule-2 (1.08 m, 1.45 m). The shortest plant height was observed in F₁ (0.86 m and 1.17 m respectively) at 6 MAP and 9 MAP. Among the interaction, S₄F₆ (3.0 m × 3.0 m with 100% of fertilizer dose as per schedule-2) recorded the highest plant height (0.75 m, 1.23 m) at 3 MAP and 6 MAP. At nine months after planting also S₄F₆ (1.60) recorded the highest plant height and it was on par with S₄F₅.

The observed increase in plant height under higher planting density may be attributed not only to competition among plants, primarily for light, but also for water and soil nutrients (Johnson and Robinson, 2000; Policarpo *et al.*, 2006) [6, 16]. Under closer spacing, overlapping canopies reduce light penetration to the leaves, thereby limiting photosynthetic efficiency and carbohydrate synthesis essential for growth (Policarpo *et al.*, 2006) [16]. As a result, intense competition for resources under high density leads to reduced shoot growth after pruning. The growth parameters were observed to be enhanced by the application of nutrients in the present investigation. This response may be attributed to the role of mineral fertilizers in enriching soil nutrient status, which in turn facilitated uptake by plant roots and supported various metabolic processes during growth (Baviskar *et al.*, 2018) [3]. These observations are consistent with the findings of Tripathy *et al.* (2015) [28] and Singh *et al.* (2016) [26].

Plant girth: Among the spacings S₁-3.0 m × 3.0 m (7.66 cm) recorded a greater plant girth and it was on par with S₂ (7.42 cm) at six months after planting. Similar trend was noted at nine months after planting also *i.e.* S₁ (12.15 cm). With respect to fertilizer application highest plant girth was recorded by F₆-100% of fertilizer dose as per schedule-2 (8.57 cm) and it was on par with F₅-100% of fertilizer dose as per schedule-1 (8.39 cm) at six months after planting. Similar trend was noted at the 9 MAP also *i.e.* F₆ (12.18 cm). Among the interaction effect, S₁F₄ (4.44 cm) recorded the highest plant girth and it was on par with S₁F₃, S₁F₁, S₁F₂, S₃F₂ and S₁ F₆ (3.94, 3.82, 3.78, 3.78 and 3.63 cm) at 3 MAP. At 6 MAP, S₁F₆ (9.30 cm) recorded the highest plant girth, and it was on par with S₁F₅, S₂F₅ (8.87 cm and 8.67 cm respectively) (Table 2). At nine months after planting also S₁F₅ (13.15 cm) recorded the highest plant girth and it was on par with S₁F₆ (13.13 cm).

Increased plant girth in higher plant spacing may be due to less competition between plants for moisture, nutrients and sunlight. Similar observations were made by Bal and Dhaliwal (2003) [2]. This response may be attributed to greater photosynthetic activity and higher chlorophyll content in the leaves of trees planted at wider distances. The absorption of nitrogen and phosphorus likely facilitated the synthesis of proteins and amino acids, contributing to the development of new tissues. Furthermore, the application of balanced nutrient doses appeared to stimulate indole-3-acetic acid (IAA) production, which promoted cell division and consequently resulted in increased stem girth.

E W Direction (cm)

The highest plant spread at E-W Direction was noticed in S₁ (35.75 cm) at 3 MAP. S₁ (84.15 cm and 120.08 cm) recorded higher plant spread at 6 MAP and 9 MAP. Results obtained from the individual effect of fertilizer application level revealed that F₃ (36.91cm) recorded the highest plant

spread at 3 MAP. Highest plant spread was recorded on F₅ (88.49 cm) at 6 MAP. However, at 9 MAP it was recorded on F₆ (120.23 cm). Among the interaction, S₁F₅ (99.33 cm) was superior and it was on par with S₁F₆ (94.33 cm) at 6 MAP and at 9 MAP S₁F₆ (136.01 cm) recorded higher plant spread (Table 3).

Greater plant spread under wider spacing may be attributed to enhanced inter-row distance, which facilitates improved light interception (Leigh, 1999) [13]. Conversely, reduced plant growth was observed under closer spacing, primarily due to reduced light penetration into the canopy and restricted apical growth, ultimately limiting overall plant vigour. Similar observations on the influence of plant spacing in guava have been reported by Pratibha *et al.* (2013) [17] and Pal and Lal (2015) [15].

N-S Direction (cm): Individual effect of spacing on plant spread revealed that planting in S₁ spacing resulted in the highest plant spread in N-S direction at 3 MAP (42.59 cm), 6 MAP (81.94 cm) and 9 MAP (102.72 cm) which were significantly different from other spacing treatments except at 9 MAP, where S₂ (99.63 cm) was on par. Plant spread was recorded to be the lowest for S₃ at 3 MAP (23.81 cm), and S₄ at 6 MAP (66.25 cm) and 9 MAP (76.18 cm). The different fertilizer doses showed significant variation where F₃ (35.66 cm) recorded higher plant spread in N-S direction at 3 MAP, F₆ (96.16 cm) recorded the highest plant spread in N-S direction at 6 MAP and F₅ (104.16 cm) recorded the highest plant spread in N-S direction at 9 MAP and this was on par with F₆ (102.30 cm). Plant spread was recorded to be the least in F₁ at 3 MAP, 6 MAP and 9 MAP (21.94 cm, 53.13 cm, and 73.25 cm). The interaction effect revealed that significant difference was observed among the treatments at 6 MAP and 9 MAP. Plant spread was the highest for S₁F₆ (105.98 cm) which was on par with S₁F₅ (100.55 cm) and S₂F₆ (99.22 cm) at 6 MAP. However, S₁F₅ (123.44 cm) recorded the highest plant spread at 9 MAP and it was on par with S₂F₅ (117.22 cm) and S₁F₆ (110.66 cm). Baviskar *et al.* (2018) [3] observed that application of nitrogen, phosphorus, and potassium has been reported to significantly enhance the vegetative growth of guava plants.

Primary branch girth (cm): Primary branch girth of the plant did not show any significant difference at different spacings at 3 MAP, 6 MAP and 9 MAP. Fertilizer dose significantly influenced primary branch girth in guava at 6 MAP and 9 MAP. 100% of fertilizer dose as per schedule-1 (F₅) registered the highest primary branch girth at 6 MAP (4.50 cm), and 9 MAP (7.21 cm), which was significantly

different from other spacing treatments except at 6 MAP where F₆ (4.39 cm) is found to be on par. The interaction level of spacing and fertilizer dose did not show any significant variation among the treatments at any observational stage (Table 5).

Secondary branch girth (cm): Spacing did not show any significant variation in the secondary branch girth at 3 MAP. S₁ (3.46 cm) spacing showed the highest secondary branch girth and it was on par with S₂ (3.45 cm) at 9 MAP. Different doses of fertilizer and interaction effect of spacing and fertilizer dose did not show any significant variation among the treatments at 3 and 9 MAP.

Pruned biomass (kg): Different levels of spacing did not express any significant difference among the treatments. Application of 100% of fertilizer dose as per schedule-2 (F₆) recorded higher pruned biomass at 6 MAP (0.16 kg) and 9 MAP (0.26 kg) and it was on par with 100% of fertilizer dose as per schedule-1 (F₅, 0.15 kg and 0.22 kg). The interaction level of spacing and fertilizer application level did not show any significant variation among the treatments at all observation stages (Table 5). In guava and other perennial fruit crops, the application of higher fertilizer doses has been shown to increase pruned biomass, primarily through the stimulation of vegetative growth. Adequate nutrient supply, particularly of nitrogen, phosphorus, and potassium, enhances cell division, shoot elongation, and lateral branching, thereby contributing to the development of a denser canopy (Sharma *et al.*, 2013) [19]. Improved nutrient availability also increases leaf area and chlorophyll content, resulting in greater photosynthetic efficiency and assimilate production, which ultimately leads to higher biomass accumulation (Kumar *et al.*, 2017) [9].

Days to shoot emergence: Spacing did not show any significant variation in the days to shoot emergence at 3 MAP and 6 MAP. S₁ (6.16) treatment resulted in the least number of days to shoot emergence at 9 MAP followed by S₂ (6.49). Fertilizer dose significantly influenced days to shoot emergence in guava i.e. emergence of primaries after first pruning and secondaries after second pruning). F₆ showed the least number of days to shoot emergence after first (6.32) and second (5.86) pruning and after second pruning F₆ was on par with F₅ (6.57). The interaction effects of spacings and fertilizer doses showed significant variation among the treatments during 9 MAP. S₁F₆ (5.04) recorded the least number of days to shoot emergence followed by S₂F₆ (5.67).

Table 1: Schedule 2:

Time of application	N (%)	P (%)	K (%)
After first pruning	40	100	20
After second pruning	30	—	30
After third pruning	30	—	50
Total	100	100	100

Table 2: Effect of spacing and fertilizer dose on plant height and plant girth at 3 MAP, 6 MAP and 9 MAP.

Treatments	Plant height (m)			Plant girth (cm)		
	3 MAP	6 MAP	9 MAP	3 MAP	6 MAP	9 MAP
Spacing level (S)						
S ₁	0.66	0.83	1.17	3.86	7.66	12.15
S ₂	0.38	0.93	1.27	2.67	7.42	11.47
S ₃	0.52	1.01	1.33	2.86	7.04	10.63
S ₄	0.65	1.08	1.42	3.40	7.03	9.77
SEm (+)	0.034	0.004	0.008	0.32	0.114	0.085
CD (0.05)	0.119	0.015	0.027	NS	0.395	0.296
Fertilizer dose (F)						
F ₁	0.54	0.86	1.17	3.24	6.14	9.71
F ₂	0.61	0.90	1.19	3.52	5.95	10.19
F ₃	0.57	0.94	1.23	3.22	7.18	10.88
F ₄	0.55	0.98	1.33	3.18	7.49	11.27
F ₅	0.47	1.01	1.41	2.97	8.39	11.80
F ₆	0.57	1.08	1.45	3.05	8.57	12.18
SEm (+)	0.021	0.003	0.009	0.14	0.127	0.059
CD (0.05)	0.062	0.010	0.028	NS	0.364	0.169
Interaction (S x F)						
S ₁ F ₁	0.75	0.76	1.14	3.82	6.49	10.51
S ₁ F ₂	0.67	0.78	1.07	3.78	6.63	11.29
S ₁ F ₃	0.59	0.80	1.04	3.94	7.32	12.2
S ₁ F ₄	0.70	0.84	1.21	4.44	7.37	12.63
S ₁ F ₅	0.65	0.86	1.26	3.55	8.87	13.15
S ₁ F ₆	0.62	0.96	1.32	3.63	9.30	13.13
S ₂ F ₁	0.30	0.86	1.16	3.45	6.67	10.16
S ₂ F ₂	0.41	0.87	1.22	3.04	6.26	10.46
S ₂ F ₃	0.46	0.89	1.23	2.7	7.09	11.60
S ₂ F ₄	0.30	0.93	1.31	1.88	7.33	11.63
S ₂ F ₅	0.29	0.98	1.33	1.97	8.67	12.61
S ₂ F ₆	0.49	1.03	1.35	2.99	8.53	12.38
S ₃ F ₁	0.49	0.90	1.16	2.68	6.23	9.50
S ₃ F ₂	0.60	0.97	1.21	3.78	5.67	9.90
S ₃ F ₃	0.66	1.01	1.27	2.89	6.83	10.49
S ₃ F ₄	0.62	1.02	1.34	3.01	7.29	10.84
S ₃ F ₅	0.32	1.03	1.46	2.69	8.03	11.16
S ₃ F ₆	0.75	1.12	1.52	2.09	8.21	11.92
S ₄ F ₁	0.60	0.92	1.22	3.01	5.20	8.70
S ₄ F ₂	0.75	1.00	1.28	3.50	5.27	9.11
S ₄ F ₃	0.57	1.05	1.38	3.35	7.50	9.24
S ₄ F ₄	0.59	1.12	1.47	3.40	7.98	10.00
S ₄ F ₅	0.63	1.18	1.58	3.68	8.01	10.27
S ₄ F ₆	0.75	1.23	1.60	3.49	8.24	11.31
SEm (+)	0.043	0.007	0.019	0.29	0.255	0.118
CD (0.05)	0.124	0.020	0.058	0.84	0.729	0.338

Table 3: Effect of spacing and fertilizer dose on E-W direction and N-S direction at 3 MAP, 6 MAP and 9 MAP.

Treatments	E W Direction (cm)			N S Direction (cm)		
	3 MAP	6 MAP	9 MAP	3 MAP	6 MAP	9 MAP
Spacing level (S)						
S ₁	35.75	84.15	120.08	42.59	81.94	102.72
S ₂	22.57	78.79	112.51	26.32	73.92	99.63
S ₃	27.16	75.37	98.90	23.81	69.74	79.27
S ₄	30.23	69.66	85.10	25.92	66.25	76.18
SEm (+)	2.420	0.956	0.178	2.910	0.768	1.941
CD (0.05)	8.40	3.309	0.617	10.07	2.657	6.718
Fertilizer dose (F)						
F ₁	17.49	63.10	87.88	21.94	53.13	73.25
F ₂	34.58	67.80	95.01	31.74	56.36	80.10
F ₃	36.91	79.66	101.38	35.66	72.33	83.69
F ₄	33.30	78.16	106.01	30.91	70.22	93.19
F ₅	23.71	88.49	114.39	27.66	89.58	104.16
F ₆	27.58	84.75	120.23	30.05	96.16	102.30
SEm (+)	3.070	0.858	0.240	2.820	1.635	2.409
CD (0.05)	8.770	2.455	0.688	8.07	4.674	6.886
Interaction (S x F)						
S ₁ F ₁	15.22	69	101.39	39.33	55.11	94.89
S ₁ F ₂	43.77	73.28	107.58	43.55	63.78	98.66
S ₁ F ₃	42.55	89	120.76	49.55	89.34	87.88
S ₁ F ₄	39.66	80	123.54	41.55	76.89	100.78
S ₁ F ₅	41.66	99.33	131.23	39.66	100.55	123.44
S ₁ F ₆	31.66	94.33	136.01	41.88	105.98	110.66
S ₂ F ₁	24.66	61.75	98.00	25.77	54.55	76.22
S ₂ F ₂	20.21	70.33	106.22	23.88	58.22	88.44
S ₂ F ₃	31.44	82.33	110.40	35.44	74.43	103.33
S ₂ F ₄	22.88	82.99	114.66	29.55	71.89	105.00
S ₂ F ₅	11.55	89.99	121.03	18.44	85.22	117.22
S ₂ F ₆	24.66	85.37	124.77	24.88	99.22	107.56
S ₃ F ₁	15.77	60.99	80.74	10.77	51.64	67.55
S ₃ F ₂	43.11	71.61	90.99	35.99	56.91	70.66
S ₃ F ₃	38.99	77.00	93.68	31.11	66.78	70.22
S ₃ F ₄	33.77	75.33	100.41	27.44	68.77	85.77
S ₃ F ₅	18.55	88.66	111.5	14.88	86.66	83.99
S ₃ F ₆	12.77	78.67	116.08	22.66	87.66	97.44
S ₄ F ₁	14.33	60.66	71.39	11.88	51.22	54.33
S ₄ F ₂	31.22	56.00	75.27	23.55	46.55	62.66
S ₄ F ₃	34.66	70.33	80.67	26.55	58.77	73.33
S ₄ F ₄	36.88	74.33	85.44	25.10	63.33	81.22
S ₄ F ₅	23.10	76.00	93.78	37.66	85.89	91.99
S ₄ F ₆	41.21	80.66	104.06	30.79	91.77	93.55
SEm (+)	6.140	1.717	0.481	5.650	3.271	4.818
CD (0.05)	NS	4.909	1.377	NS	9.349	13.772

Table 4: Effect of spacing and fertilizer dose on primary branch girth and secondary branch girth at 3 MAP, 6 MAP and 9 MAP.

Treatments	Primary branch girth (cm)			Secondary branch girth (cm)	
	3 MAP	6MAP	9MAP	3 MAP	9MAP
Spacing level (S)					
S ₁	1.92	4.25	6.69	0.56	3.46
S ₂	1.97	4.19	6.70	0.49	3.45
S ₃	1.80	3.89	6.54	0.31	3.21
S ₄	1.91	4.05	6.29	0.44	3.21
SEm (+)	0.193	0.120	0.106	0.230	0.042
CD (0.05)	NS	NS	NS	NS	0.145
Fertilizer dose (F)					
F ₁	1.98	3.75	6.08	0.15	3.29
F ₂	2.01	3.76	6.03	0.43	3.13
F ₃	1.90	4.10	6.37	0.67	3.34
F ₄	1.86	4.08	6.78	0.54	3.28
F ₅	1.94	4.50	7.21	0.50	3.45
F ₆	1.71	4.39	6.88	0.42	3.51
SEm (+)	0.127	0.139	0.119	0.160	0.097
CD (0.05)	NS	0.399	0.341	NS	NS
Interaction (S x F)					
S ₁ F ₁	1.71	3.62	6.04	0.00	3.29
S ₁ F ₂	2.21	4.21	6.01	0.45	3.18
S ₁ F ₃	2.27	4.25	6.44	0.68	3.42
S ₁ F ₄	1.80	4.15	7.09	0.47	3.26
S ₁ F ₅	1.65	4.80	7.55	0.91	3.93
S ₁ F ₆	1.88	4.51	7.05	0.83	3.70
S ₂ F ₁	2.39	3.81	6.09	0.63	3.35
S ₂ F ₂	1.88	3.58	5.92	0.39	3.09
S ₂ F ₃	1.93	4.25	6.91	0.58	3.55
S ₂ F ₄	1.93	4.57	6.6	1.13	3.47
S ₂ F ₅	2.37	4.56	7.55	0.00	3.32
S ₂ F ₆	1.32	4.40	7.12	0.22	3.90
S ₃ F ₁	1.60	3.71	6.05	0.00	3.27
S ₃ F ₂	2.11	3.47	6.08	0.68	3.05
S ₃ F ₃	1.54	3.98	6.22	0.43	3.22
S ₃ F ₄	1.92	3.46	6.89	0.20	3.18
S ₃ F ₅	1.95	4.37	7.08	0.43	3.29
S ₃ F ₆	1.69	4.35	6.94	0.16	3.26
S ₄ F ₁	2.20	3.85	6.13	0.00	3.25
S ₄ F ₂	1.83	3.81	6.11	0.20	3.20
S ₄ F ₃	1.88	3.91	5.90	0.98	3.18
S ₄ F ₄	1.82	4.16	6.53	0.36	3.23
S ₄ F ₅	1.78	4.29	6.67	0.66	3.25
S ₄ F ₆	1.97	4.29	6.4	0.47	3.18
SEm (+)	0.255	0.279	0.238	0.320	0.195
CD (0.05)	NS	NS	NS	NS	NS

Table 5: Effect of spacing and fertilizer dose on pruned biomass and days to shoot emergence at 3 MAP, 6 MAP and 9 MAP.

Treatments	Pruned biomass (kg)			Days to shoot emergence		
	3 MAP	6MAP	9MAP	3 MAP	9MAP	9MAP
Spacing level (S)						
S ₁	0.19	0.14	0.22	7.16	6.98	6.16
S ₂	0.04	0.11	0.19	7.22	6.96	6.49
S ₃	0.04	0.11	0.16	7.22	6.93	6.88
S ₄	0.05	0.10	0.13	7.27	7.16	7.45
SEm (+)	0.057	0.023	0.020	0.161	0.054	0.048
CD (0.05)	NS	NS	NS	NS	NS	0.166
Fertilizer dose (F)						
F ₁	0.04	0.07	0.10	7.58	7.78	7.56
F ₂	0.25	0.07	0.13	7.50	7.67	7.16
F ₃	0.05	0.11	0.17	7.58	7.00	6.91
F ₄	0.05	0.12	0.17	7.08	6.70	6.64
F ₅	0.03	0.15	0.22	6.75	6.57	6.32
F ₆	0.04	0.16	0.26	6.83	6.32	5.86
SEm (+)	0.080	0.012	0.028	0.266	0.124	0.040
CD (0.05)	NS	0.036	0.080	NS	0.356	0.114
Interaction (S x F)						
S ₁ F ₁	0.07	0.09	0.12	7.66	7.77	7.13
S ₁ F ₂	0.85	0.11	0.16	7.66	7.46	6.55
S ₁ F ₃	0.07	0.13	0.19	7.66	6.89	6.23
S ₁ F ₄	0.07	0.14	0.19	7.00	6.72	6.21
S ₁ F ₅	0.03	0.16	0.25	6.33	6.81	5.77
S ₁ F ₆	0.04	0.19	0.38	6.66	6.22	5.04
S ₂ F ₁	0.03	0.06	0.11	8.00	7.69	7.33
S ₂ F ₂	0.06	0.06	0.14	7.33	7.67	6.92
S ₂ F ₃	0.05	0.11	0.2	7.66	7.55	6.72
S ₂ F ₄	0.04	0.12	0.16	6.33	6.55	6.33
S ₂ F ₅	0.02	0.14	0.26	7.66	6.11	6.00
S ₂ F ₆	0.04	0.16	0.27	6.33	6.19	5.67
S ₃ F ₁	0.03	0.06	0.09	7.33	7.77	7.6
S ₃ F ₂	0.07	0.06	0.10	7.33	7.77	7.2
S ₃ F ₃	0.04	0.10	0.15	7.33	6.72	7.07
S ₃ F ₄	0.05	0.12	0.17	7.66	6.88	6.69
S ₃ F ₅	0.03	0.14	0.19	6.66	6.11	6.53
S ₃ F ₆	0.03	0.15	0.23	7.00	6.33	6.18
S ₄ F ₁	0.03	0.06	0.09	7.33	7.87	8.20
S ₄ F ₂	0.04	0.06	0.12	7.66	7.78	7.97
S ₄ F ₃	0.04	0.10	0.13	7.66	6.84	7.63
S ₄ F ₄	0.04	0.11	0.14	7.33	6.67	7.34
S ₄ F ₅	0.05	0.14	0.16	6.33	7.27	7.01
S ₄ F ₆	0.07	0.15	0.17	7.33	6.55	6.57
SEm (+)	0.161	0.025	0.056	0.532	0.249	0.080
CD (0.05)	NS	NS	NS	NS	NS	0.266

Conclusion

On the basis of experimental results, it can be concluded that the adoption of the widest plant spacing of 3.0 m × 3.0 m with application of 100% of fertilizer dose as per schedule-1 (i.e. equal splits) and 100% of fertilizer dose as per schedule-2 resulted in better growth parameters like plant girth, canopy spread (N-S direction and E-W direction), primary branch girth, secondary branch girth and pruned biomass and least number of days to shoot emergence while closer spacing of 2.0 m × 1.5 m with 100% of fertilizer dose as per schedule-2 resulted in the highest plant height in guava under HDP in the humid tropical climatic region of Kerala.

Acknowledgement

The authors are highly grateful to Kerala Agricultural University for providing the funding and facilities for carrying out the work.

Conflict of Interest

All authors declared that there is no conflict of interest

References

1. Ajithpal S, Dhaliwal GS. Influence of radiation interception and canopy temperature on fruit quality of *Psidium guajava* L. at different planting distances. *Indian J Hort.* 2004;61(2):118-121.
2. Bal JS, Dhaliwal GS. High-density planting studies in *Psidium guajava* L. *Haryana J Hort Sci.* 2003;32(1):19-20.
3. Baviskar MN, Bharad SG, Nagre PK. Effect of NPK fertilization on growth and yield of *Psidium guajava* L. under high density planting. *Int J Chem Stud.* 2018;6(3):359-362.
4. Chavan MT, Thutte AS, Kakade AR, Solanke AA. Effect of levels of N, P, K on yield and quality of *Psidium guajava* L. under high density planting. *J Pharmacogn Phytochem.* 2020;9(6):1290-1293.
5. Dwivedi V. Effect of integrated nutrient management on yield, quality and economics of *Psidium guajava* L. *Ann Plant Soil Res.* 2013;15(2):149-151.
6. Johnson PR, Robinson DM. The Tatura trellis system for high density mangoes. *Acta Hortic.* 2000;509:359-364.
7. KAU [Kerala Agricultural University]. Package of Practices Recommendations: Crops. 15th ed. Thrissur: Kerala Agricultural University; 2016. p.393.
8. Khan S, Kumar A, Sharma JR. Impact of NPK application on growth and yield of *Psidium guajava* L. cv. Hisar Safeda. *Int J Curr Microbiol App Sci.* 2018;7(7):286-290.
9. Kumar J, Prasad VM, Singh SK. Effect of nutrient management on growth and yield of *Psidium guajava* L. *J Hortic Sci Biotechnol.* 2017;92(5):529-534.
10. Kumar P, Tiwari JP, Lal S, Rajkumar. Effect of NPK levels on growth and yield of *Psidium guajava* L. *Hortic J Sci.* 2008;3(1):43-47.
11. Kumar VD, Pandey K, Anjaneyulu, Vishal N. Optimization of major nutrients for *Psidium guajava* L. yield and quality under east coastal conditions. *Indian J Hort.* 2009;66:18-21.
12. Kundu S. Effect of high density planting on growth, flowering and fruiting of *Psidium guajava* L. *Acta Hortic.* 2007;735:267-270.
13. Leigh L. Closer planting of peach trees in Goulburn Valley orchards, Tatura. *NY Fruit Q.* 1999;7:26-30.
14. Maity PK, Das BC, Kundu S. Effect of different sources of nutrients on yield and quality of *Psidium guajava* L. cv. L-49. *J Crop Weed.* 2006;2(2):17-19.
15. Pal M, Lal S. Effect of different high density planting on growth and yield of *Psidium guajava* L. cv. Pant Prabhat. *Int J Basic Appl Agric Res.* 2015;13(3):420-424.
16. Policarpo M, Talluto G, Bianco RL. Vegetative and productive responses of 'Conference' and 'Williams' pear trees planted at different in-row spacings. *Sci Hortic.* 2006;109:322-331. doi:10.1016/j.scienta.2006.06.009
17. Pratibha LS, Goswami AK. Effect of pruning and planting systems on growth, flowering, fruiting and yield of *Psidium guajava* L. cv. Sardar. *Indian J Hortic.* 2013;70(4):496-500.
18. Reddy BMC. High density planting in horticultural crops. In: First Indian Horticulture Congress, 6-9 November, New Delhi; 2004. p.36.
19. Sharma RR, Patel VB, Singh R. Nutrient management in fruit crops-a review. *Indian J Hortic.* 2013;70(1):1-15.
20. Singh G, Rajan S, Singh AK. Approaches and strategies for precision farming in *Psidium guajava* L. In: Singh HP, Singh Gorakh, Samual JC, Pathak RK, editors. Precision Farming in Horticulture. NCPAH, DAC, MoA, PFDC, CISH; 2003. p.92-113.
21. Singh A. Light interception behaviour of *Psidium guajava* L. and its effects on vegetative growth, fruit yield and quality. Ph.D. Thesis, Punjab Agricultural University, Ludhiana; 2003.
22. Singh G. Recent development in production of *Psidium guajava* L. *Acta Hortic.* 2007;735:161-173.
23. Singh G, Singh AK, Mishra D. High density planting in *Psidium guajava* L. *Acta Hortic.* 2007;735:[pp. xx-xx].
24. Singh HJ, Bal JS. Effect of planting density on tree growth, fruit yield and quality of *Psidium guajava* L. 'Sardar'. *J Res Punjab Agric Univ.* 2002;39:56-62.
25. Singh R. High density planting studies in *Psidium guajava* L. M.Sc. Thesis, Punjab Agricultural University, Ludhiana; 2006.
26. Singh TK, Mahajan G, Kumar A, Kumar P, Tiwari RK, Singh J. Growth, yield and quality of *Psidium guajava* L. as influenced by different levels of nutrients under rainfed region of Kymore plateau. *Bioscan.* 2016;11(1):275-277.
27. Tripathi A. Studies on canopy management in high density planting of *Psidium guajava* L. Ph.D. Thesis, College of Agriculture, CCSAU, Hisar, Haryana; 2018.
28. Tripathy P, Sethi K, Patnaik AK, Mukherjee SK. Nutrient management in high density cashew plantation under coastal zones of Odisha. *Int J Bio Resour Stress Manag.* 2015;6(1):93-97.