



ISSN Print: 2617-4693  
 ISSN Online: 2617-4707  
 IJABR 2023; 7(1): 39-43  
[www.biochemjournal.com](http://www.biochemjournal.com)  
 Received: 21-10-2022  
 Accepted: 29-11-2022

**Alagbe John Olujimi**  
 Department of Animal  
 Nutrition and Biochemistry,  
 Sumitra Research Institute,  
 Gujarat, India

## Bioactive compounds in ethanolic extract of *Strychnos innocua* root using gas chromatography and mass spectrometry (GC-MS)

**Alagbe John Olujimi**

DOI: <https://doi.org/10.33545/26174693.2023.v7.i1a.161>

### Abstract

Medicinal plants are of great relevance with endless pharmaceutical and therapeutic properties. They are source of wide array of secondary metabolites or bioactive compounds/phytochemicals (phenols, alkaloids, flavonoids, terpenoids, tannins, steroids and saponins) which are capable of discharging numerous biological functions (antimicrobial, antioxidant, antiviral, antifungal, anti-fibrotic, immunomodulatory, cytotoxic, antipyretic, antitumor, antihelminthic, antiprotozoal, antibacterial and so on). The use of gas chromatography and mass spectrometry analysis in identifying the bioactive compounds in ethanolic extract of *Strychnos innocua* root is a fundamental technique in quantifying the unknown samples, trace elements and contaminants leading to the discovery of novel compounds of pharmaceutical and biomedical importance. Result obtained showed that *Strychnos innocua* root extract contains 39 bioactive compounds with  $\alpha$ -Cubebene (20.09%) having the highest concentration followed by Dibutyl benzene-1,2-dicarboxylate (10.17%),  $\beta$ -Elemenone (10.02%), 4-Methoxy-2-nitroformanilide (7.21%), 1-Methyl cyclopropane methanol (5.96%), 1, 3 propanediol, 2-ethyl 2-hydroxymethyl (3.71%), Azelaic acid (2.87%), Glycidol stearate (2.85%), Chloromethyl 2-chlorodecanoate (2.83%) and  $\gamma$ -terpinene (2.56%) respectively. The remaining 29 bioactive compounds have concentrations less than 2%. It was concluded that all the compounds observed are sources of medication that can be used traditionally in the treatment of human and animal diseases.

**Keywords:** *Strychnos innocua*, phytochemicals, pharmaceuticals, therapeutics, gas chromatography

### Introduction

Medicinal plants are the major components of almost all indigenous or alternative systems of medicines. They contain phytochemicals which are safe, non-toxic and easily affordable (Singh *et al.*, 2022; Shittu and Alagbe, 2020) [9, 15]. According to WHO (1996) [39] about 80% of the population in developing countries rely on medicinal plants for the treatment of various ailments. There are over 2000 medicinal plants with high potential that are yet to be explored (Oluwafemi *et al.*, 2020) [12]. Bioactive compounds from these medicinal plants can perform an anti-inflammatory, antifungal, antiviral, antioxidant, immune-stimulatory, analgesics, antibacterial, anti-proliferative, cytotoxic and hepato-protective, antipyretic, antihelminthic, antiprotozoal, anti-depressant, anti-tumor, anti-fibrotic and hypolipidemic properties (Alagbe, 2022; Olafadehan *et al.*, 2021; Agubosi *et al.*, 2022) [4, 16, 10-11] and could also aid in the discovery of drugs (Vasquez *et al.*, 2017; Hirovani *et al.*, 1991; Muritala *et al.*, 2022) [7, 29, 3].

*Strychnos innocua* also known as Natal orange belongs to the family Loganiaceae, genus *Strychnos* and order Gentianales. The tree is found in several countries such as; Angola, Guinea, Madagascar, Malawi, South Africa, Sudan, Mali, Uganda, Malawi, Zimbabwe, Zambia, Ethiopia and some parts of India (Maghembe, 1994) [41]. The trees are found in riverine fringes, sand forest and it can grow up to 3-14 m high with simple leaves characterized by rounded emarginate or subacute apex (Hines and Eckman, 1993) [40]. Extracts from the leaves, roots and stems can be used traditionally for the treatment of snake bites, gastrointestinal, skin diseases, pneumonia and sexually transmitted infections (Al-Wathnani, 2012) [37].

**Corresponding Author:**  
**Alagbe John Olujimi**  
 Department of Animal  
 Nutrition and Biochemistry,  
 Sumitra Research Institute,  
 Gujarat, India

Previous studies have revealed that *Strychnos innocua* leaf, stem and root extract contains several bioactive compounds with antimicrobial properties and are also capable of inhibiting the activity of some bacteria and fungi such as; *Bacillus* spp, *Candida* spp, *Alternaria solani*, *Brevibacillus brevis*, *Cochliobolus lunatus*, *Escherichia coli*, *Enterobacter* spp, *Aspergillus* spp, *Fusarium* spp, *Klebsiella* spp, *Monascus ruber*, *Micrococcus luteus*, *Pseudomonas* spp, *Streptococcus* spp, *Styphlococcus* spp, *Salmonella* spp and *Shigella shiga*. Phytochemical analysis of methanolic extract from leaves and roots of *Strychnos innocua* revealed the presence of phenolic compounds which are capable of scavenging free radicals (antioxidants) (Hamisu *et al.*, 2021; Lee *et al.*, 2011; Igbal *et al.*, 2011) [36, 23, 25], preventing the risk of cardiovascular disease (Alagbe *et al.*, 2022; Alagbe, 2021) [4, 18] and performing immune-modulatory activities in animals thus encouraging food safety (Oloruntola *et al.*, 2018; Halliwell and Gutteridge, 1998) [2, 1]. The aim of this study was to evaluate the bioactive compounds in ethanolic extract of *Strychnos innocua* root using gas chromatography and mass spectrometry (GC-MS).

## Materials and Methods

### Experimental site, collection and preparation of *Strychnos innocua* ethanolic leaf and root extract

The study was performed at Sumitra Research Institute, Gujarat, India with a coastline of 1,600 Km, 23° 13'N

72°41'E. Fresh *Strychnos innocua* root was harvested from Waghai village, Saputara, India and identified at the Department of Biological Sciences, Sumitra Research Institute, Gujarat, India. The harvested roots was washed with distilled water, dried under the shade for 13 days and blended into powder form with the aid of electric blender and kept in an air tight labeled container. 100 g of grinded sample was soaked in 350 mL of 90% ethanol for 48 hours with occasional stirring thereafter samples was sieved using Whatman's No. 1 filter paper (10 cm) and stored in a sterile air tight container and stored in a cool dry place until further use.

### Gas chromatography and mass spectrometry (GC-MS) of ethanolic extract from *Strychnos innocua* root

Gas chromatography mass spectrometry (GC-MS) analysis of ethanolic extract from *Strychnos innocua* root was performed with a Varian 450 GC system (Model 1842 series, China) equipped with fused silica column and it was operated at a temperature and pressure range of 50 °C to 450 °C isothermal 1079 PTV injector and 0 to 100 psi, consisting of splitless injector with total flow of 500 mL/minutes at 10 psi, electron range of 150eV. Secondary compounds were identified with standard compounds in National Institute of Standard and Technology (NIST).

**Table 1:** Secondary metabolites in *Strychnos innocua* ethanolic root extract using GC-MS

Bioactive compounds	Area (%)	R.T (min)	Functions
Di-ethyl suberate	1.44	1.450	Antimicrobial and antioxidant
Ethyl Oleate	0.72	1.931	Antipyretic and antioxidant
Diisooctyl phthalate	0.01	2.500	Anti-depressant and antifungal
Glycidol stearate	2.85	3.444	Anitviral, hepato-protective and antioxidant
1,2 – Benzenedicarboxylic acid	1.77		Anti-microbial, anti-proliferative
Monomethyl pimelate	0.02	6.091	Antifungal
$\gamma$ -terpinene	1.10	9.435	Hepatoprotective and antifungal
4-fluoro-1-methyl-5-carboxylic acid	0.40	10.701	Anti-inflammatory, antibacterial and analgesics
3-Allyl-6-methoxyphenol	1.67	11.331	Antiprotozoal and cytotoxic
Cyclooctane	0.10	15.560	Anti-androgenic, antiviral and anti-inflammatory
Formamide	2.05	15.740	Hepato-protective, hypolipidemic, antimicrobial and antioxidant
$\alpha$ -cubebene	20.09	15.100	Antibacterial, antifungal, angelsics antipyretic and antioxidant
2,4,6 –Octatrien-1-ol	0.77	15.607	Antiviral and antioxidant
9,12-Octadecanoic acid	1.06	18.351	Cytotoxic, antioxidant, anti-inflammatory, antitumor, antifungal
$\alpha$ -longipinene	0.15	18.220	Anti-inflammatory, antioxidant, anti-depressant and antifungal
Azelaic acid	2.87	18.306	Anti-fibrotic, anti-inflammatory and hypolipidemic
Terpinen-4-ol	1.51	18.331	
1,3 propanediol, 2-ethyl 2-hydroxymethyl	3.71	18.211	Antibacterial, anti-inflammatory, antipyretic, antihelminthic and antifungal
$\gamma$ -Terpinene	2.56	19.386	Amtioxidant and anti-inflammatory
$\beta$ -Elemenone	10.02	19.931	Cytotoxic and hepato-protective
9-Octadecenoic acid	1.16	19.510	Antifungal
Torreyol- $\alpha$ -cadinol	0.83	19.259	Anitviral, hepato-protective and antioxidant
Hepatadec-3-enal	0.30	19.400	Anti-microbial, anti-proliferative, antiviral, antihelminthic and antibacterial
Ethylene diacrylate	0.50	20.209	Analgesics, antibacterial, antifungal
1-Hexyl -2 nitrocyclohexane	1.62	21.344	Anti-inflammatory, antioxidant, anti-depressant
Chloromethyl 2-chlorodecanoate	2.83	21.381	Anti-fibrotic, anti-inflammatory
5-methylhexan-2-yl hepdecyl benzene -1,2 dicarboxylate	1.10	21.100	Antifungal, angelsics antipyretic and antioxidant
1-Methyl cyclopropane methanol	5.96	22.891	Hepatoprotective and antifungal
4-Acetoxy-3-methoxystyrene	1.14	23.080	Hepato-protective, hypolipidemic, antimicrobial
4-Methoxy-2-nitroformanilide	7.21	23.300	Cytotoxic, antioxidant
$\alpha$ -Terpinolene	0.02	23.701	Antioxidant, anti-proliferative
3-deoxy-d-mannoic acid	1.20	25.186	Antioxidant, anti-proliferative, antifungal and anti-inflammatory
Dibutyl benzene -1,2 - dicarboxylate	10.17	25.901	Cytotoxic and hepato-protective

Butyl undecyl benzene-1,2- dicarboxylate	1.31	25.670	Cytotoxic, antioxidant, anti-inflammatory, antitumor, antifungal
9,15 – Octadecadienoic acid	0.06	26.801	Antibacterial, antifungal, angelsics antipyretic
β-Cyclocitral	0.01	28.009	antitumor, antifungal
α-Phellandrene	1.10	34.491	Angelsics antipyretic and antioxidant
β-Phenethylamine	0.08	38.567	Anti-bacterial
Humulene	1.12	41.340	Antimicrobial, antifungal and hypolipidemic
Total	92.59		

R.T: reaction time (minutes)

## Results and Discussion

Medicinal plants contain natural compounds or phytochemicals that are eco-friendly, safe and locally available with pharmacological properties (Musa *et al.*, 2020; Adewale *et al.*, 2021) [14, 13]. They can also be used traditionally for the treatment of various ailments such as cold, cough, gastrointestinal disease, skin disease, respiratory disease, malaria, typhoid and snake bites (Nascimento *et al.*, 2000) [42]. Bioactive compounds are mostly secondary metabolites produced by plants via subsidiary pathways and are used by plants for growth, or defense against pathogens (Okeke *et al.*, 2001; Oluwafemi *et al.*, 2021; Narayani *et al.*, 2012) [43, 45, 12]. Secondary metabolites in *Strychnos innocua* ethanolic root extract using gas chromatography and mass spectrometry (GC-MS) (Table 1) reveals that it is largely contains α-Cubebene (20.09%), Dibutyl benzene -1,2 – dicarboxylate (10.17%), β-Elementone (10.02%), 4-Methoxy-2-nitroformanilide (7.21%), 1-Methyl cyclopropane methanol (5.96%), 1,3 propanediol, 2-ethyl 2-hydroxymethyl (3.71%), Azelaic acid (2.87%), Glycidol stearate (2.85%), Chloromethyl 2-chlorodecanoate (2.83%) and γ-terpinene (2.56%) respectively. Other compounds reported were less than 2.0% however, they all have a marked therapeutic functions (anti-inflammatory, antiviral, antifungal, antioxidant, hypolipidemic, angelsics, anti-pyretic, cytotoxic, antitumor and anti-depressant activities) (Okeke *et al.*) [43]. The GC-MS component analyzed in this study is in consonance with the findings of Hamisu *et al.* (2021) [36] but contrary to the reports of Hoet *et al.* (2006) [30]. These dissimilarity can be ascribed to processing or extraction procedures employed, parts of plant used, species, geographical location, age of plant as well as method of harvesting (Omokore and Alagbe, 2019; Hoet *et al.*, 2006) [17, 30]. The presence of phytochemicals in *Strychnos innocua* ethanolic root extract reveals that it has the ability to scavenge toxic chemicals in the body, inhibit the activities of pathogenic bacteria in the gut of animals, thus enhancing the absorption of nutrient as well as enhances the activities of enzymes (Oluwafemi *et al.*, 2021; Narayani *et al.*, 2012) [12, 45].

## Conclusion

Medicinal plants have so several health benefits due to the presence of phytochemicals (alkaloids, flavonoids, phenols, terpenoids, saponins, tannins and steroids). Analyzing the bioactive components in *Strychnos innocua* ethanolic root extract will unleash some of the potential pharmaceutical properties in the plant. Adopting the use of gas chromatography and mass spectrometry will further aid in identifying and quantifying unknown samples, unknown contaminants, trace elements and gases.

## References

- Halliwell B, Gutteridge JMC. Free radicals in biology and medicine. Oxford University Press, Oxford, UK; c1998.

- Oloruntola OD, Agbede JO, Ayodele SO, Adeyeye SA, Agbede JO. Performance, haemato-biochemical indices and antioxidant status of growing rabbits fed on diets supplemented with *Mucuna pruriens* leaf meal. World Rabbit Science. 2018;26:277-285.
- Shittu MD, Alagbe JO, Ojebiyi OO, Ojediran TK, Rafiu TA. Growth performance and haematological and serum biochemical parameters of broiler chickens given varied concentrations of *Polyalthia longifolia* leaf extract in place of conventional antibiotics. Animal Science and Genetics. 2022;18(2):57-71.
- Olujimi AJ, Muhammad RS, Daniel SM, Christiana OO. Effect of *Trichilia monadelpha* stem bark extract on the fatty acid composition of rabbit's thigh meat. Journal of Environmental Issues and Climate Change. 2022;1(1):63-71.
- Ruth TN, Anita RL, Loveness KN, Vincenzo F, Ruud V. Local processing and nutritional composition of indigenous fruits: the case of monkey orange (*Strychnos* spp.) from Southern Africa. Food Research. 2017;33(2):123-142.
- Alagbe JO, Shittu MD, Tanimomo Babatunde K. Influence of *Anogeissus leo carpus* stem bark on the fatty acid composition in meat of broiler chickens. European Journal of Life Safety and Stability. 2022;14(22):13-22.
- Vasquez-Ocmin P, Cojean S, Rengifo C, Suyyagh-Albouz S, Guerra CAA, Pomel S, *et al.* Antiprotozoal activity of medicinal plants used by Iquitos-Nauta road communities in Loreto (Peru). Journal of Ethno Pharmacology. 2017;4(61):1-9.
- Alagbe JO. Use of medicinal plants as a panacea to poultry production and food security: A review. International Journal of Economy and Innovation. 2022;22:1-12.
- Sharma S, John AO, Xing L, Ram S, Amita K. Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectroemetry. International Journal of Agriculture and Animal Production. 2022;2(6):18-27.
- Agubosi OCP, James A, Alagbe JO. Influence of dietary inclusion of Sunflower (*Helianthus annus*) oil on growth performance and oxidative status of broiler chicks. Central Asian Journal of Medical and Natural Sciences. 2022;2(7):187-195.
- Agubosi OCP, Soliu MB, Alagbe JO. Effect of dietary inclusion levels of *Moringa oleifera* oil on the growth performance and nutrient retention of broiler starter chicks. Central Asian Journal of Theoretical and Applied Sciences. 2022;3(3):30-39.
- Oluwafemi RA, Omolade LA, Adetope AS, Alagbe JO. Effects of dietary inclusion of ginger (*Zingiber officinale*) and garlic (*Allium sativum*) oil on carcass characteristics and sensory evaluation of broiler

- chicken. Texas Journal of Multidisciplinary Studies. 2021;2(11):180-188.
13. Adewale AO, Alagbe JO, Adeoye Adekemi O. Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phyto-genic Feed Additive in Growing Rabbit Diets: Haematology and serum biochemical indices. International Journal of Orange Technologies. 2021;3(3):1-12.
  14. Musa B, Alagbe JO, Betty AM, Omokore EA. Growth performance, caeca microbial population and immune response of broiler chicks fed aqueous extract of *Balanites aegyptiaca* and *Alchornea cordifolia* stem bark mixture. United Journal for Research and Technology. 2020;2(2):13-21.
  15. Shittu MD, Alagbe JO. Phyto-nutritional profiles of broom weed (*Sida acuta*) leaf extract. International Journal of Integrated Education. 2020;3(11):119-124.
  16. Olafadehan OA, Oluwafemi RA, Alagbe JO. Performance, haemato-biochemical parameters of broiler chicks administered Rolfe (*Daniellia oliveri*) leaf extract as an antibiotic alternative. Advances in Research and Reviews. 2020;1:4.
  17. Omokore EO, Alagbe JO. Efficacy of dried *Phyllanthus Amarus* leaf meal as an herbal feed additive on the growth performance, haematology and serum biochemistry of growing rabbits. International Journal of Academic Research and Development. 2019;4(3) 97-104.
  18. Alagbe JO. Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phyto-genic Feed Additive in Growing Rabbit Diets: Growth Performance and Caecal Microbial Population. Concept in Dairy and Veterinary Sciences; c2021, 4(2).
  19. Agubosi OCP, Wika BK, Alagbe JO. Effect of dietary inclusion of Sunflower (*Helianthus annuus*) oil on the growth performance of broiler finisher chickens. European Journal of Modern Medicine and Practice. 2022;2(5):1-10.
  20. Alagbe JO. *Prosopis africana* stem bark as an alternative to antibiotic feed additives in broiler chicks diets: Performance and Carcass characteristics. Journal of Multidimensional Research and Reviews. 2021;2(1):64-77.
  21. Orwa CA, Mutua KR, Jamnadass R, Anthony S. Agroforestry database: A tree reference. 2009. Retrieved from <http://www.worldagroforestry.org/sites/treedbs/treedata bases. asp>
  22. Alagbe JO. *Daniellia oliveri* leaf extracts as an alternative to antibiotic feed additives in broiler chicken diets: Meat Quality and Fatty acid composition. Indonesian Journal of Innovation and Applied Sciences. 2021;1(3):177-186.
  23. Lee SW, Wendy W, Julius YFS, Desy FS. Characterization of antimicrobial, antioxidant, anticancer properties and chemical composition of *Michelia champaca* seed and fower extracts. Stamford Journal of Pharmaceutical Science. 2011;4(1):19-24.
  24. Alagbe JO. Chemical evaluation of proximate, vitamin and amino acid profile of leaf, stem bark and roots of *Indigofera tinctoria*. International Journal on Integrated Education. 2020;3(10):150-157.
  25. Igbal H, Moneebur RK, Riazullah ZM, Naeem KFA, Zahoor U. Phytochemical screening and antimicrobial activities of selected medicinal plants of Khyber Pakhtunkhwa, Pakistan. African Journal of Pharmacology. 2011;5(6):746-750.
  26. Alagbe JO. Caecal Microbial Population of Growing Grass Cutters (*Thyronorm Swinderianus*) Fed *Phyllanthus Amarus* and *Piliostigma Thonningii* Leaf Meal Mixture as Partial Replacement for Soya Bean Meal. Concept of Dairy and Veterinary Sciences. 2020;3(5):350-355.
  27. Sa'Ayinzat F, Bawa E, Ogwu D, Ayo J. Hesperidin-Sources, chemistry, extraction, measurement and biologic effects on reproduction in animals: A review. International Journal of Veterinary Sciences and Animal Husbandry. 2021;6(4):1-8.
  28. Shivakumara KN. Review on friedel-crafts acylation of benzene derivatives using various catalytic systems. Int. J Adv. Chem. Res. 2021;3(1):25-31. DOI: 10.33545/26646781.2021.v3.i1a.32
  29. Hirotani H, Ohigashi M, Kobayashi K, Koshimizu K, Takahashi E. Inactivation of T5 phage by cis-vaccenic acid, an antiviral substance from *Rhodopseudomonas capsulate*, and by unsaturated fatty acids and related alcohols. FEMS Microbiological Letters. 1991;77(1):13-17.
  30. Hoet S, Stevigny C, Herent M, Quetin-Leclercq J. Antitrypanosomal compounds from the leaf essential oil of *Strychnos spinosa*. Planta Medicine. 2006;72:480-482.
  31. Alagbe JO, Habiba Z, Adedeji OM, Bamigboye S, Agbonika D. Influence of *Juniperus thurifera* root extract on the nutrient digestibility and caecal microbial count of growing rabbits. Web of Synergy: International Interdisciplinary Research Journal. 2022;1(1):5-17.
  32. Oluwafemi RA, Agubosi OCP, Alagbe JO. Proximate, minerals, vitamins and amino acid composition of *Prosopis africana* (*African mesquite*) seed oil. Asian Journal of Advances in Research. 2021;11(1):21-27.
  33. Oluwafemi RA, Uankhoba IP, Alagbe JO. Effects of turmeric oil as a dietary supplements on the growth performance and carcass characteristics of broiler chicken. International Journal of Orange Technologies. 2021;3(4):1-9.
  34. Oluwafemi RA, Uankhoba IP, Alagbe JO. Effects of turmeric oil as a dietary supplement on the haematology and serum biochemical indices of broiler chickens. Bioinformatics and Proteomics Open Access Journal. 2021;5(1):000138.
  35. Oluwafemi RA, Daniel SE, Alagbe JO. Haematology and serum biochemical indices of broiler chicks fed different inclusion levels of ginger (*Zingiber officinale*) and garlic (*Allium sativum*) oil mixture. International Journal of Discoveries and Innovations in Applied Sciences. 2021;1(4): 20-26.
  36. Ibrahim H, Uttu AJ, Sallau MS, Risikat O, Agbeke Iyun L. Gas chromatography–mass spectrometry (GC–MS) analysis of ethyl acetate root bark extract of *Strychnos innocua* (Delile). Beni-Suef University Journal of Basic and Applied Sciences. 2021;10:65.
  37. Al-Wathnani H, Ismet A, Tahmaz RR, Al-Dayel TH, Bakir MA. Bioactivity of natural compounds isolated from cyanobacteria and green algae against human pathogenic bacteria and yeast. Journal of Medicinal Plants Research. 2012;6(18):3425-3433.



38. Asuzu CU, Nwosu MO. Studies of the wood of some Nigeria alkaloid-rich *Strychnos* species. *Journal of Horticulture*. 2020;12(2):57-62.
39. WHO. Annex II. Guidelines for the assessment of herbal medicines (WHO Technical Report Series No. 863), Geneva; c1996.
40. Hines DA, Eckman K. Indigenous multipurpose trees for Tanzania: uses and economic benefits to the people. Cultural survival Canada and Development Services Foundation of Tanzania; c1993.
41. Maghembe JA. Germination studies on seed of fruit trees indigenous to Malawi. *Forest Ecology and Management*. 1994;64(2-3):111-125.
42. Nascimento GGF, Lacatelli J, Freitas PC, Silva GL. Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. *Brazil Journal Microbiology*. 2000;31(4):886-891.
43. Okeke MI, Iroegbu CU, Eze EN, Okoli AS, Esimone CO. Evaluation of extracts of the roots of *Landolphia owerience* for antibacterial activity. *Journal Ethnopharmacology*. 2001;78(2-3):119-27.
44. Trease GE, Evans WC. *Pharmacognasy* WB. Scandars Company Ltd. London. 1989;14:269-300.
45. Narayani M, Johnson M, Sivaraman A, Janakiraman N. Phytochemical and Antibacterial Studies on *Jatropha curcas* L. *Journal of Chemical and Pharmaceutical Research*. 2012;4(5):2639-2642.
46. Parekh J, Chands S. *In vitro* antibacterial activity of the crude methanol extract of *Woodfordia fruticosa* Kurz. Flower (Lythraceae), *Brazil Journal of Microbiology*. 2007;38:204-207.
47. Hongxiang S, Cuirong S, Yuanjian P. Cytotoxic activity and constituents of the volatile oil from the roots of *Patrinia scabra* Bunge. *Chemical Biodiversity*. 2005;2(10):1351-1357.
48. Alagbe JO. Gas chromatography and mass spectroscopy of *Juniperus phoenicea* stem bark extract and its influence on the haemato-biochemical values of growing rabbits. *British Scientific Periodical*. 2022;1(1):18-33.