International Journal of Advanced Biochemistry Research

ISSN Print: 2617-4693
ISSN Online: 2617-4707
IJABR 2018; 2(2): 35-42
www.biochemjournal.com
Received: 25-05-2018
Accepted: 27-06-2018

Mohy Eldin Abd El Fattah
Chemistry Department,
Faculty of Science, Suez Canal
University, Cairo, Egypt

**Mohamed Ramadan
Abdelgawad**
Biological Applications
Department, Nuclear Research
Centre, Atomic Energy
Authority, Cairo, Egypt

**Basher Abd Elghfar El
Boughdady**
Chemistry Department,
Faculty of Science, Suez Canal
University, Cairo, Egypt

The protective role of Epigallocatechin gallate (EGCG) for renal damage by suppressing oxidative stress in rats induced by aluminum oxide nanoparticles

Mohy Eldin Abd El Fattah, Mohamed Ramadan Abdelgawad and Basher Abd Elghfar El Boughdady

Abstract

The aim of this study was to evaluate the possible Protective role of EGCG on Kidney Functions and Oxidative Stress in rats induced by Aluminum oxide nanoparticles. Eight groups of rats were used; Group1, control. Group2, received $\text{Al}_2\text{O}_3\text{NPS}$ alone in a dose 50 mg/kg b.w i.p. Group 3, received Epigallocatechin gallate alone in a dose (5 mg/kg b.w. i.v.). Group 4, received Epigallocatechin gallate alone in a dose (10 mg/kg b.w. i.v.). Group 5, received $\text{Al}_2\text{O}_3\text{NPS}$ followed by simultaneous administration of Epigallocatechin gallate in a dose (5 mg/kg b.w. i.v.). Group 6, received $\text{Al}_2\text{O}_3\text{NPS}$ followed by simultaneous administration of Epigallocatechin gallate in a dose (10 mg/kg b.w. i.v.). There was a highly significant elevation in serum of AST, ALT, ALP and GGT and while, a significant decrease in albumin and total protein level but no a significant change in globulin and A/G ratio in rats treated with $\text{Al}_2\text{O}_3\text{-NPs}$ as compared to normal control rats. There was a highly significant elevation in serum of creatinine, urea and uric acid in rats treated with $\text{Al}_2\text{O}_3\text{-NPs}$ compared to normal control rats. The concentrations of creatinine, urea and uric acid were significantly decreased in rats treated with Exelon or Epigallocatechin gallate; 5 mg/kg and 10 mg/kg treated with $\text{Al}_2\text{O}_3\text{-NPs}$ when compared to $\text{Al}_2\text{O}_3\text{-NPs}$ -treated rats. Also, the present study showed a highly significant decreased in GSH concentration, SOD activity and CAT activity but significantly increased in MDA level in blood for rats treated with $\text{Al}_2\text{O}_3\text{-NPs}$ compared to normal control rats. There was a highly significant increased GSH concentration, SOD and CAT activities while the level of MDA was decreased in rats treated with Exelon or Epigallocatechin gallate; 5 mg/kg and 10 mg/kg treated with $\text{Al}_2\text{O}_3\text{-NPs}$ when compared to $\text{Al}_2\text{O}_3\text{-NPs}$ – treated rats.

Keywords: Epigallocatechin gallate, (EGCG) suppressing oxidative

Introduction

Polyphenols are natural substances that are present in drinks obtained from plants, vegetables, and fruits, such as tea, and olive oil. The largest group of polyphenols are flavonoids which mainly divided into glycosylated derivative of anthocyanidin, anthocyanins, present in colorful flowers, fruits and anthoxantins. Anthoxantins are colorless compounds further divided into several categories including flavonols flavones, falvanols, flavans, and is flavones (Butterfield *et al.*, 2002). Aromatic ring presnt in flavonoids is reduced to a heterocyclic ring then attached to a second aromatic ring. Antioxidant activity due to abundant phenolic hydroxyl groups on the aromatic ring, and the 3-OH is essential for the iron chelating activity of these compounds (Van Acker *et al.*, 1996). The importance of polyphenolic flavonoids in improving cell resistance to oxidative stress goes beyond simple scavenging and is most important for pathologies in which oxidative stress plays an important role. Numerous studies in the last 10 years have shown that polyphenols prevent or reduce the harmful effects of free radicals derived from oxygen associated with several chronic and stress - related human and animal diseases *in vitro* and *in vivo*. Oxidative stress is due to reactive oxygen species (ROS) generation and inflammation play a vital role in scientific disorders as arteriosclerosis, neurodegenerative disorder, cancer, ischemia-reperfusion injury and stroke (M. E. Götz *et al.*, 2001). Green tea contains polyphenols, including flavonoids, flavanols, flavandiols and phenolic acids, which can account for approximately 30% of the dry weight.

Correspondence
Mohy Eldin Abd El Fattah
Chemistry Department,
Faculty of Science, Suez Canal
University, Cairo, Egypt

The majority of green tea polyphenols are flavonols commonly referred to as catechins. In green tea, four types of catechins are mainly detected: Epicatechin (EC), Epicatechin-3-gallate (ECG), Epigallocatechin (EGC) and Epigallocatechin-3-gallate (EGCG). Due to differences in origin and growing conditions, the amount of catechins differs in green tea leaves (Khokhar & Magnusdottir, 2002). Nanoparticles (NPs) may be defined as materials that have at least one dimension less than 100 nm (Balasubramanyam *et al.*, 2009). They are desirable for industrial and healthcare applications because of their unique chemical, mechanical and biological properties (Oberdörster, 2005). Because of their unique chemical, mechanical, and biological properties they are desirable for industrial and health care applications [8].

Materials and Methods

Animals

In the present study, adult male albino rats (120 ± 20 g) from the animal house of Faculty of Veterinary Medicine Suez Canal University, Egypt, were used as experimental animals. The rats were grouped in special cages with six animals per cage and maintained under our laboratory conditions; temperature (23±2), with dark and light cycle (12/12h). Standard pellet diet and water were allowed free access *ad libitum*. The rats were adapted to laboratory conditions for 7 days before starting of experiment. All procedures of experiment were performed between 8-11 a.m.

Chemicals

EGCG (M.W: 476.39, CAS Number: 989-51-5, Catalog No.: 4524, Batch No.: 2 B/189017) was purchased from Tocris Bioscience / clinilab company (4,160St. El-Etehad Square Riham Tower El-Maadi, Cairo, Egypt). Aluminum oxide nanoparticles (Al₂O₃NPS) from Egyptian Atomic Energy Authority, Inshas Science City. Chemicals used for analytical reagent grade were obtained from EGY-CHEM for lab technology, Badr city, Egypt and Biodiagnostic Company, Dokki, Giza, Egypt.

Experimental design

The rats were randomly divided into 6 groups: Group 1; received 1ml saline 0.9% orally daily throughout the experiment and served as normal control group. Group (2); received Al₂O₃NPS alone in a dose 50 mg/ kg b.w intraperitoneally (i.p), three times a week for three weeks, served as positive control group [9]. Group 3; received Epigallocatechin gallate alone in a dose (5 mg/kg b.w. i.v.) every day for five weeks. Group 4; received Epigallocatechin gallate alone in a dose (10 mg/kg b.w. i.v.) every day for five weeks [10]. Group 5; received Al₂O₃NPS in a dose 50 mg/kg b.w intraperitoneally (i.p), three times a week for three weeks followed by simultaneous administration of Epigallocatechin gallate in a dose (5 mg/kg b.w. i.v.) every day for five weeks. Group 6; received Al₂O₃NPS in a dose (50 mg/kg b.w intraperitoneally (i.p), three times a week for three weeks followed by simultaneous administration of Epigallocatechin gallate in a dose (10 mg/kg b.w. i.v.) every day for five weeks.

Biochemical Assays

A. Kidney Functions

1. Determination of serum creatinine

The concentration of creatinine was determined by fixed rate colorimetric method as described by Henry (1974)

using available commercial kit which was purchased from a local chemical company.

Procedure

- 1.0 ml of working solution (R1: 1 volume + R2: 1 volume) was added to all tubes.
- 100 µl of sample, 100 µl of standard were added to sample tube and standard tube.
- All tubes were mixed well, the initial absorbance (A1) of the standard and specimen were read at 492 nm, and then after exactly 2 minutes, the absorbance (A2) of both standard and specimen were read again.

Calculation

[A2 - A1 = Aspecimen or Astandard]

$$\text{Creatinine concentration (mg/dl)} = \frac{(A) \text{ Sample}}{(A) \text{ Standard}} \times 2.0$$

Where, 2 is the standard creatinine concentration.

2. Determination of serum urea

The level of urea was determined by a colorimetric method as described by Chaney *et al.* (1962) using available commercial kit which was purchased from a local chemical company.

Procedure

- 50 µl of reagent 2 and 1.0 ml of reagent 3 were added to Blank tube.
- 50 µl of reagent 2, 1.0 ml of reagent 3 and 10 µl of sample were added to sample tube.
- 50 µl of reagent 2, 1.0 ml of reagent 3 and 10 µl of reagent 1 standard were added to standard tube, all tubes were mixed well
- After incubation for 3 min at 37°C. and then 200 µl of reagent 4 was added to all tubes. The absorbance of sample and standard tubes was read at λ 578 nm against blank.

Calculation

The level of urea in sample was calculated using the following equation:

$$\text{Urea (mg/dl)} = \frac{(A) \text{ Sample}}{(A) \text{ Standard}} \times 50, \text{ Where, 50 is the standard urea concentration.}$$

3. Determination of serum uric acid

The concentration of uric acid was determined by a colorimetric method as described by Trinder (1969) using available commercial kit which was purchased from a local chemical company.

Procedure

- 20 µl of distilled water was added to blank tube, 20 µl of sample was added to sample tube and 20 µl of standard was added standard tube
- 1.0 ml of reagent 2 was added to all tubes.
- All tubes were mixed well.
- After incubation for 5 min. at 37°C. The absorbance of standard and sample tubes was read at 500 nm against blank.

Calculation

The level of uric acid was calculated using the following equation

$$\text{Uric acid concentration (mg/dl)} = \frac{A_{\text{specimen}}}{A_{\text{standard}}} \times 6$$

Where, 6.0 is the standard uric acid concentration.

B. Assessment of oxidative stress biomarkers

Lipid Peroxidation: Lipid peroxidation was estimated by measuring thiobarbituric acid reactive substances (TBARS) and was expressed in terms of malondialdehyde (MDA) content by a colorimetric method according to Satoh (1978).

Antioxidant Enzymes: Superoxide dismutase activity was determined according to the method of Nishikimi *et al.*, (1972). The method is based on the ability of SOD enzyme to inhibit the phenazine methosulphate-mediated reduction of nitro blue tetrazolium dye (NTB). Briefly, 0.05 mL sample was mixed with 1.0 mL buffer (pH 8.5), 0.1 mL nitro blue tetrazolium (NBT), and 0.1 mL NADH. The reaction was initiated by adding 0.01 mL phenazine methosulphate (PMs), and then increase in absorbance was read at 560 nm for five minutes.

Catalase activity was determined according to the method of Aebi (1985). The method is based on the decomposition of H₂O₂ by catalase. The sample containing catalase is incubated in the presence of a known concentration of H₂O₂. After incubation for exactly one minute, the reaction is quenched with sodium azide. The amount of H₂O₂ remaining in the reaction mixture is then determined by the

oxidative coupling reaction of 4-aminophenazone (4-aminoantipyrene, AAP) and 3,5-dichloro-2-hydroxybenzenesulfonic acid (DHBS) in the presence of H₂O₂ and catalyzed by horseradish peroxidase (HRP). The resulting quinoneimine dye (N-(4-antipyril)-3-chloro-5-sulfonate-p-benzoquinonemonoimine) is measured at 510 nm.

Erythrocyte GSH was measured following the method of Beutler, (1984). The method was based on the ability of the -SH group to reduce 5,5-dithiobis,2-nitrobenzoic acid (DTNB) and form a yellow coloured anionic product whose OD is measured at 412 nm. Concentration of GSH is expressed in milligram per millilitre packed RBCs and was determined from standard plot.

Statistical analysis

The result values were expressed as means $\pm$ standard error (SE) for 6 rats in each group. Tabulation and graphics were designed using Microsoft Excel XP software. Data were statistically analyzed using Statistical Package for Social Science (SPSS) version 19, software. One-way analysis of variance (ANOVA) test was performed to statistical analysis for determining the statistical significant differences between means of different groups. Data were considered instatistically significant when the P values were > 0.05 .

Results

• Effect of Epigallocatechin gallate on control rats Effect on blood antioxidant parameters

As shown in Table (1) and Figures (1,2,3 and 4), there was no significant variation in GSH, CAT, MDA and SOD activities compared to normal control group.

Table 1: Blood Glutathione, antioxidant enzymes and serum malodialdehyde in control, and normal rats treated with Epigallocatechingallate (n=6).

Groups / Parameters	GSH(mg/dl)	SOD(U/ml)	CAT(U/L)	MDA (nmol/ml)
Control Range (n=6)	21.99 $\pm$ 0.62 ^a (20.31-23.85)	318.57 $\pm$ 8.91 ^{bc} (301-358.1)	346 $\pm$ 20.3 ^a (316-445)	4.33 $\pm$ 0.22 ^a (3.7-5.1)
Exelon Range (n=6)	23.28 $\pm$ 1.4 ^a (16.65-25.62)	355.68 $\pm$ 10.33 ^a (320.3-386.8)	350 $\pm$ 12.35 ^a (320-399)	4.8 $\pm$ 0.23 ^a (4.17-5.5)
% Change compared to control	5.87	11.64	1.16	10.8
EGCG (10 mg)	24.50 $\pm$ 1.17 ^a (21.21-28.7)	353.82 $\pm$ 12.05 ^{ab} (310-385)	350 $\pm$ 13.33 ^a (289-380)	4.38 $\pm$ 0.24 ^a (4.0-5.4)
Range (n=6) % Change compared to control	11.41	11.07	1.16	1.15

Data presented as Mean $\pm$ SEM

Means have the same letters considered insignificant (P >0.05).

Fig 1: Mean Blood Glutathione (GSH) concentration (mg/dL) in normal control and Epigallocatechingallate groups in normal rats.

Fig 2: Mean Blood Superoxide dismutase (SOD) concentration (U/ml) in normal control and Epigallocatechingallate groups in normal rats.

Fig 3: Mean serum Catalase (CAT) Activity (U/L) in normal control and Epigallocatechingallate groups in normal

Fig 4: Mean serum MDA concentration (nmol/ml) normal control and Epigallocatechingallate groups in normal rats.

• Effect of Epigallocatechin gallate on rats treated with Al₂O₃-NPs

Effect on blood antioxidant enzymes

Effect on glutathione (GSH) contents

Results given in Table (2) and graphically illustrated in Figure (5) showed that the intraperitoneal injection of Al₂O₃-NPs in the previously mentioned dose and period to normal rats induced significantly decreased in GSH content when compared with the normal control group.

Rats treated with Exelon with Al₂O₃-NPs administration in the previously mentioned dose and period had significant increase in GSH content compared to Al₂O₃-NPs -treated rats. The percentage of increase was 44.00% compared with the Al₂O₃-NPs -treated rats. GSH contents in rats treated with Epigallocatechin (5 mg) and (10 mg) with Al₂O₃-NPs administration were increased by 71.18% and 55.86% respectively compared to Al₂O₃-NPs -treated rats.

Effect on superoxide dismutase (SOD) activity

Results in Table (2) and Figure (6) showed that the intraperitoneal injection of Al₂O₃-NPs in the previously mentioned dose and period to normal rats induced decreased in SOD activity by 23.75% compared to normal control rats. Rats treated with Epigallocatechin gallate (5 mg and 10 mg) with Al₂O₃-NPs administration in the previously mentioned dose and period had increased in SOD activity compared to Al₂O₃-NPs -treated rats. SOD activities of these rats restored to the values of normal group (309 ± 23.18, 345 ± 8.71 vs. 318.57 ± 8.91) respectively.

Effect on catalase (CAT) activity

Results in Table (2) and Figure (7) showed that the intraperitoneal injection of Al₂O₃-NPs in the previously mentioned dose and period to normal rats induced significantly decreased in CAT activity by 36.13% compared to normal control rats. Al₂O₃-NPS -treated rates with (5 mg and 10 mg) in the previously mentioned dose had significant increase in CAT activity compared to Al₂O₃-NPs -treated rats. CAT activities of these rats succeeded to restore the activities of CAT to normal values (399 ± 8.96, 405.7 ± 18.38 vs. 346 ± 20.3) respectively.

Effect on lipid peroxidation (MDA) level

Results in Table (2) and Figure (8) showed that intraperitoneal injection of AL₂O₃-NPS in the previously mentioned dose and period to normal rats induced significantly increased in malodialdehyde (MDA) level by 177.3% compared to normal control group. AL₂O₃-NPS -treated rates with Epigallocatechin gallate (5 mg and 10 mg) in the previously mentioned dose and period had a significant decrease in MDA level compared to AL₂O₃-NPS-treated rats. MDA levels of these rats returned nearly to the normal values.

Table 2: Blood Glutathione, antioxidant enzymes and serum malodialdehyde in control, AL₂O₃-NPS-treated rats, and AL₂O₃-NPS-treated rats and supplemented with Epigallocatechingallate.

Groups / Parameters	GSH(mg/dl)	SOD(U/ml)	CAT(U/L)	MDA(nmol/ml)
Control	21.99 ± 0.62 ^b	318.57 ± 8.91 ^{ab}	346 ± 20.3 ^a	4.33 ± 0.22 ^c
Range (n=6)	(20.31-23.85)	(301-358.1)	(316-445)	(3.7-5.1)
AL ₂ O ₃ -NPS	14.75 ± 1.23 ^c	242.92 ± 3.79 ^b	221 ± 6.31 ^b	12.01 ± 0.48 ^a
Range (n=6)	(10.24-18.6)	(231-255)	(205-245)	(10.5-13.8)
%Change compared to control	-32.92	-23.75	-36.13	177.3
EGCG (5 mg)	25.25 ± 0.56 ^a	309 ± 23.18 ^{ab}	399 ± 8.96 ^a	9.75 ± 4.16 ^b
Range (n=6)	(22.26-26.43)	(196.8-346.5)	(368-422.1)	(8.5-10.9)
%Change compared to control	14.82	-2.95	15.32	125.1
%Change compared to Al ₂ O ₃ -NPs	71.18	27.28	80.54	-18.8
EGCG (10 mg)	22.99 ± 0.82 ^{ab}	345 ± 8.71 ^a	405.7 ± 18.38 ^a	9.4 ± 0.39 ^b
Range (n=6)	(19.72-25.3)	(315.8-366.2)	(317.9-439)	(8.0-10.6)
%Change compared to control	-4.35	8.3	17.25	117.1
%Change compared to Al ₂ O ₃ -NPs	55.86	42	83.57	-21.73

Data presented as Mean ± SEM

Means have the same letters considered insignificant (P>0.05).

Fig 5: Mean Blood Glutathione (GSH) concentration (mg/dL) in normal control and different groups of AL2O3-NPS -treated rats.

Fig 7: Mean serum CAT activity (U/L) in normal control and different groups of AL2O3-NPS -treated rats.

Fig 6: Mean Blood SOD concentration (U/ml) in normal control and different groups of AL2O3-NPS -treated rats.

Fig 8: Mean serum MDA concentration (nmol/ml) in normal control and different groups of AL2O3-NPS -treated rats.

• **Effect of Exelon and Epigallocatechin gallate on kidney functions in control rats**

In Table (3) and Figure (9, 10 and 11), There was no variation in creatinine and urea of normal rats treated with Epigallocatechin gallate (5 mg/kg and 10 mg/kg) compared to normal control rats.

Table 3: Renal functions in control, and normal rats treated with Epigallocatechin gallate

Groups / parameters	Creatinine(mg/dl)	Urea(mg/dl)	Uric acid(mg/dl)
Control Range (n=6)	0.65± 0.006 ^a (0.63–0.67)	39 ± 2.6 ^a (28–45)	1.26 ± 0.02 ^b (1.2–1.35)
EGCG (5 mg) Range (n=6)	0.64 ± 0.25 ^a (0.61–0.77)	35 ± 1.54 ^a (31–39)	1.54 ± 0.09 ^{a b} (1.25–1.83)
%Change compared to control	22.4	-10.25	22.2
EGCG (10 mg) Range (n=6)	0.62 ± 0.31 ^a (0.55–0.75)	32 ± 1.07 ^a (29–36)	1.35 ± 0.1 ^b (1.0–1.68)
%Change compared to control	-4.6	-17.9	7.1

Data presented as Mean ± SEM

Means have the same letters considered insignificant ($P > 0.05$).

Fig 9: Mean serum creatinine concentration (mg/dL) in normal control and Epigallocatechin gallate groups in normal rats

Fig 10: Mean serum urea concentration (mg/dL) in normal control and Epigallocatechin gallate groups in normal rats.

Fig 11: Mean serum Uric acid concentration (mg/dL) in normal control and Epigallocatechin gallate groups in normal rats.

• **Effect of Exelon and Epigallocatechin gallate on kidney functions functions in rats treated with Al₂O₃-NPs**

Results given in Table (4) and graphically illustrated in Figures (12,13 and 14) showed that the intraperitoneal injection of Al₂O₃-NPs in the previously mentioned dose and period to normal rats induced significantly increased compared to normal control rats.

The concentration of creatinine, urea and uric acid were significantly decreased in Epigallocatechin (5 mg/kg or 10 mg/kg) treated with Al₂O₃-NPs administration for each when compared with the Al₂O₃-NPs group. The levels of creatinine, urea and uric acid of these rats returned nearly to the levels of control group in case of Epigallocatechin (10 mg/kg) than Epigallocatechin (5 mg/kg) treated with Al₂O₃-NPs.

Fig 13: Mean serum urea concentration (mg/dL) in normal control and different groups of AL₂O₃-NPS -treated rats.

Fig 12: Mean serum creatinine concentration (mg/dL) in normal control and different groups of AL₂O₃-NPS -treated rats.

Fig 14: Mean serum Uric acid concentration (mg/dL) in normal control and different groups of AL₂O₃-NPS -treated rats.

Table 6: Renal functions in control, AL₂O₃-NPS -treated rats, and AL₂O₃-NPS -treated rats and supplemented with Epigallocatechin gallate

Group / Parameters	Creatinine(mg/dl)	Urea(mg/dl)	Uric acid(mg/dl)
Control	0.65 ± 0.006 ^a	39 ± 2.6 ^{c,d}	1.26 ± 0.02 ^c
Range (n=6)	(0.63–0.67)	(28–45)	(1.2–1.35)
AL ₂ O ₃ -NPS	0.9 ± 0.012 ^b	60 ± 0.73 ^a	3.3 ± 0.11 ^a
Range (n=6)	(0.88–0.95)	(58–63)	(2.9–3.62)
%Change compared to control	38.4	53.9	161.9
AL ₂ O ₃ -NPS + EGCG (5 mg)	0.7 ± 0.028 ^a	59 ± 3.53 ^a	1.63 ± 0.15 ^{b,c}
Range (n=6)	(0.61–0.79)	(49–72)	(1.19–2.19)
%Change compared to control	7.7	51.2	29.3
%Change compared to AL ₂ O ₃ -NPs	-22.2	-1.7	-50.6
AL ₂ O ₃ -NPS + EGCG (10 mg)	0.6 ± 0.016 ^a	43 ± 3.7 ^{b,d}	1.4 ± 0.16 ^{b,c}
Range (n=6)	(0.56–0.66)	(30–55)	(1.09–1.99)
%Change compared to control	-7.7	10.2	11.1
%Change compared to AL ₂ O ₃ -NPs	-33.3	-28.3	-57.5

Data presented as Mean ± SEM

Means have the same letters considered insignificant (P>0.05).

Discussion

Green tea is one of human consumption's most popular drinks. Epidemiological studies have shown that green tea consumption is associated with a reduced risk of many chronic diseases, including cardiovascular diseases, diabetes and various cancers [15-18]. Green tea's health benefits can be attributed primarily to catechins, its main bioactive components. Five major catechins have been identified in green tea, including catechin (C), epicatechin (EC), epigallocatechin (EGC), epicatechin gallate (ECG) and Epigallocatechin gallate (EGCG) [19, 20].

Catechins appear to be capable of producing and scavenging free radicals and showing their beneficial effects by combining the two mechanisms [21, 22]. Catechins' antioxidant efficacy is exercised by (1) direct mechanisms - scavenging chelating ROS, metal ions; and (2) indirect mechanisms - inducing antioxidant enzymes, inhibiting pro-oxidant enzymes, and producing detoxification enzymes and antioxidant enzymes in Phase II [23]. All catechins and their diastereoisomers have common chemical structures - phenolic hydroxyl groups that can stabilize free radicals [24]. Phenolic hydroxyl groups of catechins can react in a termination reaction with reactive oxygen and reactive nitrogen species that breaks the cycle of new radical's generation. Catechins donate one phenolic OH group electron, thereby reducing free radicals and maintaining stability through the resonance of the resulting aroxyl radicals [25, 26]. The number of molecule hydroxyl groups is positively correlated with the antioxidant activity of phenolic compounds [27]. Catechins' relative efficacy hierarchy as radical scavengers is EGCG > ECG > EGC > EC [27-29].

Epigallocatechin gallate is the most potent antioxidant compound in green tea, along with its most abundant polyphenol [30]. Due to its structure of phenol rings, Epigallocatechin gallate has a powerful antioxidant activity, acts as scavengers and free radical electron traps [31, 32], Preventing the formation of reactive oxygen species and reducing oxidative stress damage [33].

The kidney is a complex organ made up of well-defined components that work in a highly coordinated way. It has been shown that a number of drugs, chemicals and heavy metals alter its structure and function, but acute and chronic intoxication has been shown to cause nephropathy with different levels of severity ranging from tubular dysfunction to acute renal failure (Barbier *et al.*, 2005). The Nephrotoxicity is of critical concern when selecting new drug candidates during the early stage of drug development because of its unique metabolism; the kidney is an important target of the toxicity of drugs, xenobiotics and oxidative stress (Uehara *et al.*, 2007).

In the present study, intraperitoneal injection of Al₂O₃-NPs to normal rats induced nephrotoxicity and renal dysfunction as evidenced by significantly increased creatinine, urea and uric acid compared to normal control rats, which suggested possible renal toxicity of alumina NPs, these results agree with (Yang *et al.*, 2012).

In the present study, the concentration of creatinine, urea and uric acid were significantly decreased in rats treated with EGCG. This effect elicited by EGCG might be due to its potent antioxidant property, as antioxidants have been reported previously for their ability to alleviate oxidative damage (Babu *et al.*, 2008; Widlansky *et al.*, 2007).

Conclusion

The present study elucidated the beneficial effects of green tea Epigallocatechin gallate evident by improvement of hepatic, renal and hematological parameters. So, our present work recommends the usage of green tea to overcome the abnormal changes in body functions. Since, green tea has been consumed over long periods without any known side effects, its possible role as an adjunct therapeutic agent against the hepatotoxicity.

References

- Weinreb O *et al.* Neurological mechanisms of green tea polyphenols in Alzheimer's and Parkinson's diseases. *The Journal of nutritional biochemistry*. 2004; 15(9):506-516.
- Choi Y-T *et al.* The green tea polyphenol (-)-Epigallocatechin gallate attenuates β -amyloid-induced neurotoxicity in cultured hippocampal neurons. *Life sciences*. 2001; 70(5):603-614.
- Nakagawa K, Miyazawa T. Absorption and distribution of tea catechin, (-)-epigallocatechin-3-gallate, in the rat. *Journal of nutritional science and vitaminology*. 1997; 43(6):679-684.
- Dulloo AG *et al.* Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24-h energy expenditure and fat oxidation in humans. *The American journal of clinical nutrition*. 1999; 70(6):1040-1045.
- Zhu QY *et al.* Regeneration of α -tocopherol in human low-density lipoprotein by green tea catechin. *Journal of agricultural and food chemistry*. 1999; 47(5):2020-2025.
- Shixian Q *et al.* Green tea extract thermogenesis-induced weight loss by Epigallocatechin gallate inhibition of catechol-O-methyltransferase. *Journal of medicinal food*. 2006; 9(4):451-458.
- Balasubramanyam A *et al.* *In vivo* genotoxicity assessment of aluminium oxide nanomaterials in rat peripheral blood cells using the comet assay and micronucleus test. *Mutagenesis*. 2009; 24(3):245-251.
- Oberdörster G, Oberdörster E, Oberdörster J. Nanotoxicology: an emerging discipline evolving from studies of ultrafine particles. *Environmental health perspectives*. 2005; 113(7):823.
- Shah SA *et al.* Nanoscale-alumina induces oxidative stress and accelerates amyloid beta (A β) production in ICR female mice. *Nanoscale*. 2015; 7(37):15225-15237.
- Rasoolijazi H *et al.* The beneficial effect of (-)-epigallocatechin-3-gallate in an experimental model of Alzheimer's disease in rat: A behavioral analysis. *Iranian Biomedical Journal*. 2007; 11(4):237-243.
- Tietze W. *Fundamentals of Clinical Chemistry* 2nd Ed WB Saunders Co., Philadelphia, 1982.
- Gornall AG, Bardawill CJ, David MM. Determination of serum proteins by means of the biuret reaction. *Journal of biological chemistry*. 1949; 177(2):751-766.
- Dumas B, Biggs H. *Standard methods of clinical chemistry*. Academic Press, Chicago. 1972; 7:175-189.
- Buttarelli M, Plebani M. Automated blood cell counts: state of the art. *American journal of clinical pathology*. 2008; 130(1):104-116.

15. Gao YT *et al.* Reduced risk of esophageal cancer associated with green tea consumption. JNCI: Journal of the National Cancer Institute. 1994; 86(11):855-858.
16. Iso H *et al.* The relationship between green tea and total caffeine intake and risk for self-reported type 2 diabetes among Japanese adults. Annals of Internal Medicine. 2006; 144(8):554-562.
17. Kuriyama S *et al.* Green tea consumption and mortality due to cardiovascular disease, cancer, and all causes in Japan: the Ohsaki study. Jama. 2006; 296(10):1255-1265.
18. Kapoor S. Re: Green tea consumption and prostate cancer risk in Japanese men: a prospective study. American journal of epidemiology. 2008; 168(1):119.
19. Seeram NP *et al.* Catechin and caffeine content of green tea dietary supplements and correlation with antioxidant capacity. Journal of Agricultural and Food Chemistry. 2006; 54(5):1599-1603.
20. Chen Q, Guo Z, Zhao J. Identification of green tea's (*Camellia sinensis* (L.)) quality level according to measurement of main catechins and caffeine contents by HPLC and support vector classification pattern recognition. Journal of Pharmaceutical and Biomedical Analysis. 2008; 48(5):1321-1325.
21. Oliveira-Marques V *et al.* Modulation of NF- κ B-Dependent Gene Expression by H₂O₂: A Major Role for a Simple Chemical Process in a Complex Biological Response. Antioxidants & redox signaling. 2009. 11(9):2043-2053.
22. Valko M *et al.* Free radicals and antioxidants in normal physiological functions and human disease. The international journal of biochemistry & cell biology. 2007; 39(1):44-84.
23. Youn HS *et al.* Suppression of MyD88-and TRIF-dependent signaling pathways of Toll-like receptor by (-)-epigallocatechin-3-gallate, a polyphenol component of green tea. Biochemical pharmacology. 2006; 72(7):850-859.
24. Fraga CG *et al.* Basic biochemical mechanisms behind the health benefits of polyphenols. Molecular aspects of medicine. 2010; 31(6):435-445.
25. Fan F-Y, Sang L-X, Jiang M. Catechins and their therapeutic benefits to inflammatory bowel disease. Molecules. 2017; 22(3):484.
26. Bors W *et al.* Flavonoids as antioxidants: Determination of radical-scavenging efficiencies, in Methods in enzymology. Elsevier. 1990, 343-355.
27. Rice-evans CA *et al.* The relative antioxidant activities of plant-derived polyphenolic flavonoids. Free radical research. 1995; 22(4):375-383.
28. Intra J, Kuo S-M. Physiological levels of tea catechins increase cellular lipid antioxidant activity of vitamin C and vitamin E in human intestinal caco-2 cells. Chemico-biological interactions. 2007; 169(2):91-99.
29. Fujisawa S, Kadoma Y. Comparative study of the alkyl and peroxy radical scavenging activities of polyphenols. Chemosphere. 2006; 62(1):71-79.
30. Sutherland BA, Rahman RM, Appleton I. Mechanisms of action of green tea catechins, with a focus on ischemia-induced neurodegeneration. The Journal of nutritional biochemistry. 2006; 17(5):291-306.
31. Rice-Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationships of flavonoids and phenolic acids. Free radical biology and medicine. 1996; 20(7):933-956.
32. Chung JE *et al.* Amplification of antioxidant activity of catechin by polycondensation with acetaldehyde. Biomacromolecules. 2004; 5(1):113-118.
33. Tipoe GL *et al.* Green tea polyphenols as an anti-oxidant and anti-inflammatory agent for cardiovascular protection. Cardiovascular & Haematological Disorders-Drug Targets (Formerly Current Drug Targets-Cardiovascular & Hematological Disorders). 2007; 7(2):135-144.
34. Hall C, Hall E. Human relations in education. Routledge, 2003.
35. Spencer A *et al.* Aluminium deposition in liver and kidney following acute intravenous administration of aluminium chloride or citrate in conscious rats. Human & experimental toxicology. 1995; 14(10):787-794.
36. Klein GL *et al.* Altered glycine and taurine conjugation of bile acids following aluminum administration to rats. Journal of pediatric gastroenterology and nutrition. 1989; 9(3):361-364.
37. HM O *et al.* Aluminium toxicity in rats: The role of tannic acid as antioxidant. Ass. Univ. Bull. Environ. Res. 2003; 6(2).
38. Morsy GM, Abou El-Ala KS, Ali AA. Studies on fate and toxicity of nanoalumina in male albino rats: oxidative stress in the brain, liver and kidney. Toxicology and industrial health. 2016; 32(2):200-214.
39. Yang S-T *et al.* Bioavailability and preliminary toxicity evaluations of alumina nanoparticles *in vivo* after oral exposure. Toxicology Research. 2012; 1(1):69-74.
40. Kuriakose GC, Kurup MG. Hepatoprotective effect of Spirulina lonar on paracetamol induced liver damage in rats. Asian J Exp Biol Sci. 2010; 1(3):614-623.
41. Pari L, Suresh A. Effect of grape (*Vitis vinifera* L.) leaf extract on alcohol induced oxidative stress in rats. Food and chemical toxicology. 2008; 46(5):1627-1634.
42. Babu A, Pon V, Liu D. Green tea catechins and cardiovascular health: an update. Current medicinal chemistry. 2008; 15(18):1840-1850.
43. Widlansky ME *et al.* Acute EGCG supplementation reverses endothelial dysfunction in patients with coronary artery disease. Journal of the American College of Nutrition. 2007; 26(2):95-102.
44. Thangapandiyan S, Miltonprabu S. Epigallocatechin gallate effectively ameliorates fluoride-induced oxidative stress and DNA damage in the liver of rats. Canadian journal of physiology and pharmacology. 2013; 91(7):528-537.
45. Mahieu S *et al.* Aluminum toxicity. Hematological effects. Toxicology letters. 2000; 111(3):235-242.
46. Wright JS, Johnson ER, DiLabio GA. Predicting the activity of phenolic antioxidants: theoretical method, analysis of substituent effects, and application to major families of antioxidants. Journal of the American Chemical Society. 2001; 123(6):1173-1183.