




International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
 ISSN Online: 2617-4707
 IJABR 2021; 5(1): 06-08
www.biochemjournal.com
 Received: 03-11-2020
 Accepted: 26-12-2020

Dr. Vittal BG
 Professor and Head,
 Department of Biochemistry,
 Hassan Institute of Medical
 Sciences, Hassan, Karnataka,
 India

Dr. Abhijith D
 Senior Resident,
 Department of Biochemistry,
 JSS Medical College,
 JSSAHER, Mysore,
 Karnataka, India

Corresponding Author:
Dr. Abhijith D
 Senior Resident,
 Department of Biochemistry,
 JSS Medical College,
 JSSAHER, Mysore,
 Karnataka, India

Fasting and non-fasting lipid profile: A comparative study

Dr. Vittal BG and Dr. Abhijith D

DOI: <https://doi.org/10.33545/26174693.2021.v5.i1a.56>

Abstract

Sedentary lifestyle and lack of physical activity predisposes to Hyperlipidaemia which is a modifiable risk factor for coronary artery disease. The deranged lipid profile is also seen in patients of hypertension and diabetes mellitus which are among the most common non communicable diseases seen mainly in developed and some developing countries. Dyslipidemia depending upon the severity will be monitored by Lipid profile usually done in the fasting state and based on the differential derangement of lipid levels, treated accordingly. Most of the nations have switched on to non-fasting lipid profiles owing to its convenience and feasibility of testing and also probably considering the fact that the body is exposed to non-fasting lipid levels for most time in 24 hours of the day. This pilot study tried to compare and study the fasting and non-fasting lipid profile components by doing an observational cross-sectional study on 50 patients attending the Hassan Institute of Medical Sciences laboratory services, Hassan for regular health checkups. There was no statistically significant difference between total cholesterol, HDL cholesterol and LDL cholesterol in the fasting and postprandial state, whereas triglycerides and VLDL showed a significant statistical difference with a p-value less than 0.05. Our study concludes that non-fasting lipid profile is better both technically and operationally, with better diagnostic and prognostic capability with respect to monitoring and treatment of dyslipidaemias as compared to fasting lipid profile.

Keywords: Fasting, non-fasting, lipid profile

Introduction

Lipid profile is one of the most commonly performed blood investigations encompassing Total cholesterol (TC), High Density Lipoprotein cholesterol (HDL), Low Density Lipoprotein cholesterol (LDL), Very low density lipoproteins (VLDL), and Triglycerides. The preferred sample for this investigation is fasting serum with 8-12 hours fasting^[1, 2]. Patients on their first visit are usually advised by the doctors to visit them again on a different day with fasting lipid profile laboratory reports which is most of the times inconvenient to doctors, patients as well as laboratories. From the perspective of the patient, a random, non-fasting lipid profile is practical compared with a fasting lipid profile because it does not interfere with the patient's normal life and allows going to the laboratory for the blood draw at any time of the day^[3]. From the perspective of the laboratory, a fasting lipid profile is thought to be superior to a non-fasting one because it can be standardized and leads to slightly less variation in the measured analytes^[5].

Fasting for >8 h, as required for lipid profiles, normally only occurs a few hours before breakfast. By contrast, the non-fasting state predominates most of the 24 hour cycle and better captures atherogenic lipoprotein levels^[3]. With the developing knowledge base it is known that the body is exposed to lipids round the clock and fasting lipid profile values represents only the fraction that the body is exposed for 3-4 hours and during the rest of the times the body is exposed to non-fasting lipids. So, non-fasting lipid profile values ideally should be more indicative of the status of health or disease^[3, 4]. Not many studies have been conducted in India to study the non-fasting lipid values in our population, which accounts for 1/6th population of the world. This study intends to find the difference between fasting and non-fasting lipid profiles.

Materials and Methods

An observational cross sectional study was done on patients attending the Hassan Institute

Of Medical Sciences laboratory services, Hassan for a regular health checkup with the approval of the institutional ethical committee on a pilot basis. After obtaining informed consent from the patients, the blood samples given by the patients in the fasting and postprandial state were analysed for Lipid Profile. Type of sampling employed was a sampling of convenience and age group of patients was between 30 to 70 years. Out of 50 patients recruited for the study, 20 were males and 50 were females. Fasting and postprandial blood samples obtained from the patients using standard venepuncture techniques were analysed for total cholesterol, triglycerides and HDL using ERBA EM 360 auto analyser. VLDL and LDL were calculated using Friedewald formula. Triglycerides/5 gives the VLDL value and TC-[HDL + VLDL] gives the value of LDL. The units for all the parameters of Lipid profile is mg/dL. The data collected was analysed by using graph pad prism version 9. The measured parameters were expressed as Mean \pm Standard deviation. The parameters were compared using paired 't' test at the 5 % level of significance. p-value <0.05 was considered statistically significant. Paired "t-test" was used because we measured lipid profile before and after food.

Results

A total of 50 patients of age between 30 to 70 years were involved in this study. The results obtained along with the graphs are as given below:

Table 1: Table showing the clinical parameters and their results

Parameters	Fasting Sample	Non Fasting state	p-value
Total cholesterol	217 \pm 49	214 \pm 51	0.27
Triglycerides	240 \pm 92	254 \pm 91	0.0002
HDL cholesterol	52 \pm 14	51 \pm 15	0.4285
VLDL cholesterol	48 \pm 18	51 \pm 18	0.0003
LDL cholesterol	117 \pm 46	112 \pm 48	0.0948




Fig 1: Comparison between total cholesterol v/s post prandial cholesterol and fasting total cholesterol

The mean Total cholesterol level in the fasting sample is 217 with a standard deviation of 49, the mean Total cholesterol level in non-fasting sample is 214 with a standard deviation of 51 with p-value of 0.27.




Fig 2: Comparison between triglyceride level in mg/dL v/s post prandial triglycerides and fasting triglycerides

The mean Triglyceride level in the fasting sample is 240 with a standard deviation of 92, the mean Triglyceride level

in non fasting sample is 254 with a standard deviation of 91 with a p-value of 0.0002.




Fig 3: Comparison between HDL level in mg/dL v/s post prandial HDL and fasting HDL

The mean HDL level in the fasting sample is 52 with a standard deviation of 14, the mean HDL level in non-fasting sample

is 51 with a standard deviation of 15 with p-value of 0.425.




Fig 4: Comparison between VLDL level in mg/dL v/s post prandial VLDL and fasting VLDL

The mean VLDL level in the fasting sample is 48 with a standard deviation of 18, the mean VLDL level in non-fasting sample is 51 with a standard deviation of 18 with p-value of 0.0003.




Fig 5: Comparison between LDL level in mg/dL v/s post prandial LDL and Fasting LDL

The mean LDL level in the fasting sample is 117 with a standard deviation of 46, the mean LDL level in non-fasting sample is 112 with a standard deviation of 48 with p-value of 0.0948.

Discussion

This pilot study showed that there is no statistically significant difference between total cholesterol, HDL cholesterol and LDL cholesterol in the fasting and postprandial state indicating that it should not make any difference whether the blood sample is collected in the fasting state or non-fasting state for these three parameters.

A statistically significant difference was noted in fasting and post prandial levels of triglycerides and VLDL cholesterol, indicating that fasting and non-fasting samples cannot be used interchangeably for these parameters. In our study, VLDL and LDL being calculated parameters; mainly depend on Triglycerides, Total cholesterol and HDL. Direct measurement of these two parameters could have added more strength to the statistical significance. Also, as per the Friedewald formula, validity of calculated VLDL and LDL is not good whenever the triglyceride level will be more than 400 mg/DL because of which patients with Triglyceride level more than 400 mg/dl in fasting or non-fasting state have not been included in the study.

Lynn E. Eberly, Jeremiah Stamler, James D. Neaton, *et al.* In their study published in 2003 state that fasting and non-fasting triglycerides are similarly predictive of nonfatal and fatal coronary artery disease [CAD] [6]. Robert H. Eckel in his article published in 2014 states that VLDL predicts coronary artery disease better as compared to Triglycerides [7]. Wang H and Eckel R.H in their study published in 2009 state that fasting VLDL will usually miss out the VLDL remnants which are the produced from Chylomicrons by the action of lipoprotein lipase [8]. From the above studies, it can be inferred that fasting and non-fasting triglycerides do not matter pertaining to prediction of CAD and VLDL predicts

CAD better in non-fasting state.

Due to its convenience and better clinical value, many organizations across the globe have accepted the use of non-fasting lipid profile. The Danish Society for Clinical Chemistry recommended nationwide use of non-fasting lipid profiles in 2009, followed by a similar endorsement in 2011 by the American Heart Association and, in 2014, by the U.K. National Institute for Health and Care Excellence guidelines and guidelines from Veterans Affairs and U.S. Department of Defense. This was followed by similar recommendations in 2016 by the European Atherosclerosis Society [3].

Conclusion

Irrespective of statistical significance, non-fasting lipid profile is better both technically and operationally with respect to convenience and feasibility and more advisable as our body is exposed to non-fasting state for more duration [3] as fasting state persists only for a short period of time after we wake up and even just a bed coffee or tea will further reduce the body's exposure time to fasting lipid levels. A well designed study with a large number of subjects, random sample instead of postprandial sample and direct estimation of all the parameters of lipid profile would add on more credibility and validity for correlating the fasting and non-fasting lipid profiles.

References

1. Rifai N, Warnick GR. Lipids, lipoproteins, apolipoproteins, and other cardiovascular risk factors. In: Burtis CA, Ashwood ER, Bruns DE, editors. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. 4th Edition. Philadelphia, PA: Elsevier Saunders; c2006, p. 903-982.
2. Simundic AM, Cornes M, Grankvist K, Lippi G, Nybo M. Standardization of collection requirements for fasting samples: for the Working Group on Preanalytical Phase (WG-PA) of the European Federation of Clinical Chemistry and Laboratory Medicine (EFLM). Clin Chim Acta. 2014;432:33-7.
3. Nordestgaard BG. A Test in Context: Lipid Profile, Fasting Versus Nonfasting. J Am Coll Cardiol. 2017;70(13):1637-1646
4. Nordestgaard BG, Langsted A, Mora S, *et al.* Fasting is not routinely required for determination of a lipid profile: clinical and laboratory implications, including flagging at desirable concentration cut-points: A joint consensus statement from the European Atherosclerosis Society and European Federation of Clinical Chemistry and Laboratory Medicine. EUR Heart J. 2016;37(25):1944-58.
5. Lorenzo Lozano MC, Cosmen Sanchez A, Belinchón Torres PM, Prieto Menchero S, Pineda - Tenor D. Is fasting necessary for lipid profile determinations? Some considerations from the perspective of the clinical laboratory. Clin Chem Lab Med. 2017;55(9):e187-8.
6. Lynn Eberly E, Jeremiah Stamler, James Neaton D, *et al.* Relation of Triglyceride Levels, Fasting and Nonfasting, to Fatal and Nonfatal Coronary Heart Disease. Arch Intern Med. 2003;163(9):1077-1083
7. Robert Eckel H. What is it about very low density lipoproteins (VLDL) and cardiovascular disease in patients with type 2 diabetes mellitus: Is it the triglycerides or the cholesterol? Atherosclerosis. 2014;237(1):138-139.
8. Wang H, Eckel RH. Lipoprotein lipase: from gene to obesity. Am J Physiol Endocrinol Metab. 2009;297(2):E271-88.