

ISSN Print: 2617-4693 ISSN Online: 2617-4707 IJABR 2024; SP-8(3): 471-475 www.biochemjournal.com Received: 10-01-2024 Accepted: 13-02-2024

Iqra Nazir

Department of Aquaculture, GBPUAT, Pantnagar, Uttarakhand, India

Mohd Danish Department of FRM, GBPUAT, Pantnagar, Uttarakhand, India

Sauliheen Qadri Division of FRM, SKAUST-K, Srinagar, Jammu and Kashmir, India

Tabinda Shabir Yaswi Department of Fish Processing Technology, KUFOS, Panangad, Kochi, Kerala, India

Corresponding Author: Mohd Danish Department of FRM, GBPUAT, Pantnagar, Uttarakhand, India

Assessment of biochemical parameters (Total serum protein, albumin and globulin) for immunity enhancement in fingerlings of *Cyprinus carpio haematopterus*

Iqra Nazir, Mohd Danish, Sauliheen Qadri and Tabinda Shabir Yaswi

DOI: https://doi.org/10.33545/26174693.2024.v8.i3Sf.793

Abstract

In the current study, the immunostimulatory effects of chitosan, vitamin C, *Curcuma longa, Tinospora cordifolia, Withania somnifera*, and *Allium sativum* on *Cyprinus carpio haematopterus* fingerlings were assessed. A total of 25 fish per treatment group (T₀, T₁, T₂, and T₃) were randomly assigned to receive fingerlings weighing 30.0 ± 2 g each in duplicate. A combination of rice bran, deoiled mustard oil cake, deoiled soybean cake, and vitamin-mineral mixture was used to create four isoproteinous experimental diets. The phytobiotics, Vitamin C and Chitosan were incorporated into diet. Per kilogramme of feed, diet D1 included 2.5 g each of garlic, giloy, turmeric, and ashwagandha whereas the diet D₂ had control feed + (2.5 g of each plant) + 25 mg Vitamin C per kg feed. For every kilogram of feed, diet D3 included 2.5 grams of each plant and 2.5 grams of chitosan. T₀ group fishes were fed with D₀ diet, T₁ with D₁, T₂ with D₂ and T₃ with D₃ @ 5% body weight per day for 122 days. All treatment groups fed the experimental meal showed a significant (p<0.05) increase in the levels of total blood protein, albumin, globulin, A/G ratio, and ALT between the pre-challenged and post-challenged groups. The findings show that the tested phytobiotics, in combination with chitosan and vitamin C, have excellent immunostimulatory effects and have no negative effects on the health of *Cyprinus carpio haematopterus* in a culture system.

Keywords: Total serum protein, albumin, globulin and immunity

Introduction

Asia contributes about 88.91% to the world fish production according to FAO (2016) ^[5]. Carps contribute to the majority of aquaculture production worldwide, accounting for over 70% of total productivity. These comprise of Indian major carps i.e, Catla (*Calta catla*), Rohu (*Labeo rohita*) and Mrigala (*Cirrhinus mrigala*); Chinese carps viz. Silver carp (*Hypophthalmicthys molitrix*), Grass carp (*Ctenopharyngodon idella*), Common carp (*Cyprinius carpio*) and Big head carp (*Hypophthalmicthys nobilis*). Among these Grass carp (*Ctenopharyngodon idella*) is the highest cultivated carp species contributing about 5.02 mmt, silver carp (*Hypophthalmicthys molitrix*) ranks 3rd contributing about 4.18 mmt and common carp (*Cyprinius carpio*) ranks 4th contributing 3.79 mmt to the world aquaculture production (FAO, 2015)^[4]. Aquaculture is a significant industry that is growing annually on a global scale. The Indian fisheries is also contributing to the national food security of the country. India ranks second globally in terms of both total fish production and freshwater fish production. In India, fisheries sector provides employment to about 14.49 million people for livelihood security.

Experimental site and climate

The experiment was conducted in the tarai region of Uttarakhand at the Wet Lab of the College of Fisheries, G. B. Pant University of Agriculture and Technology, Pantnagar, Distt. Udham Singh Nagar. The experiment ran from March 1 until June 30, 2018.

Fish collection and maintenance

At the Fish Seed Hatchery, College of Fisheries, G. B. Pant University of Agriculture and Technology, Pantnagar, three hundred fingerlings of healthy, disease-free *Cyprinus carpio haematopterus* were obtained. Their average body weight was 30.00 ± 2 g. The fingerlings were kept in indoor captive settings in aerated water for two days while they acclimated in cemented tanks.

Treatment combination

T₀ - D₀ Control diet

 T_1 - (D₁) with control feed + 2.5 g of garlic + 2.5 g of giloy + 2.5 g of turmeric + 2.5 g of ashwagandha per kg feed T_2 - (D₂) with control feed + (2.5 g of each of four plants) +

25 mg Vitamin C per kg feed

 T_3 - Diet (D₃) with control feed + (2.5 g of each plant four plants) + 2.5 g Chitosan per kg feed

Feeding was done @ 5% body weight daily. The total quantity of the daily diet was given in two installments, once at 9.00 am and then at 5.00 pm.

Experimental design

The entire experiment was carried out under controlled circumstances at the G. B. Pant University of Agriculture and Technology, Pantnagar, in the Wet Lab of the College of Fisheries. In this experiment, twelve equal-sized FRP tanks with a diameter of 1.07 meters and a depth of 0.82 meters were employed in four groups of three duplicates $(4T_3R=12)$. About 600 liters of water were kept in each tank. Every day, the water was aerated. There were twenty-five Amur carp fingerlings in each tank. Fish in the control treatment T_1 were given D_1 diet, T_2 fishes with D_2 diet and T_3 fishes with D_3 diet. The experiment lasted for 122 days. The growth parameters were analyzed every 15^{th} day. Biochemical analysis was done at 0 day, pre challenge, post challenge and withdrawal day.

Biochemical analysis

Total serum protein

Total serum protein was calculated by using following formula

Total serum protein $(g/dl) = (Absorbance test / Absorbance of standard) \times 6.5$

Albumin

The albumin was calculated by using following formula

Albumin (g/dl) = (Absorbance of test / Absorbance of standard) $\times 4$

Globulins

The globulin was calculated by using following formula

Globulins = Total serum protein – Albumin

Albumin-globulin ratio [A/G]

Albsumin- globulin ratio =

The albumin-globulin ratio was calculated by using following formula

Serum albumin (g/dl)

Serum Alanine aminotransferase (ALT)

The general formula for converting absorbance change into International Units (IU) of activity is: Activity of ALT = Δ Abs./min. x 1768 at 37 °C (IU/L)

Bacterial strain and challenge study

After 102 days of feeding trial, blood samples were collected from treatment and control fish. Suspension of Aeromonas hydrophila (designation RTMCX1) was prepared in PBS from freshly grown culture for the challenge experiment from diagnostic bacteriology laboratory, ICAR- DCFR, Bhimtal and maintained at 4 °C in the Department of Aquaculture, G.B Pant University of Agriculture and technology, Pantnagar. Ten fishes from each treatment were challenged intraperitoneally with 10 µl of bacterial suspension $(1.5 \times 10^8 \text{ CFU/ml})$ on day 92. Adequate care was taken to avoid injury when the strain came into contact with A. hydrophila. All exposed samples were returned to their respective containers and monitored for their reaction against the injected bacterial strain. Mortality was observed for 10 days. After 102 days of feeding the fish with the D_1 , D_2 and D_3 diet, the fishes in the all groups were then fed with control diet for 15 days. After that blood was taken for biochemical analyses.

Statistical analysis

The experimental data were subjected to a two-way analysis of variance (ANOVA) using SPSS (Statistical Package for Social Sciences 2006 version 16.0). The individual mean differences were compared using Duncan's multiple range test. When p<0.05 was reached, the values were deemed significant.

Results and Discussion Total serum protein (TSP)

The observations on total serum proteins in different treatment groups have been depicted in Figure1. The total serum protein (TSP) of amur carp at 0th day varied between 1.88±0.53 to 1.99±0.061 g/dl. The total serum protein of amur carp in pre challenge fishes varied between 2.45±0.006 to 2.76±0.069 g/dl among treatments. In pre challenged group, the maximum value for TSP was recorded as 2.76±0.069 g/dl (T₃). In case of post challenged fishes, the value ranged between 4.94 ± 0.09 to 6.92 ± 0.019 . Highest TSP was obtained in T_3 (6.92±0.019 g/dl). Between T_0 , T_1 , T_2 , and T_3 on the 0th day and pre-challenge, as well as between the pre-challenge and post-challenge groups fed experimental diets, the value of TSP differed considerably (p < 0.005). Abdel-Tawwab *et al.* (2010) ^[1] discovered that when tilapia were fed green tea and subsequently challenged with Aeromonas hydrophila, there was an increase in total serum protein, albumin, and globulin. Harikrishnan et al. (2012)^[7] also noticed that their was an increase in total serum protein when the fish were fed with 1% or 2% of chitosan diets. The results are in agreement with Siwicki (1989) ^[14] who found that after feeding carp fish with β glucan and chitosan their was an increase in the level of total protein content. The values of TSP for the withdrawal period varied between 2.42±0.006 to 2.97±0.169.

Albumin

The observations on albumin in different treatment groups have been depicted in (figure2). The albumin level of Amur carp at 0th day varied between 0.59 ± 0.051 to 0.63 ± 0.532 g/dl. The albumin level of Amur carp in pre challenged

fishes varied between 0.82±0.01 to 0.97±0.03 g/dl among treatment. In pre challenged group, the maximum value for albumin was recorded as 0.97 ± 0.03 g/dl (T₃). The value for fish that had been post- challenged ranged from 1.66±0.01 to 3.29±0.04 g/dl. The T₃ treatment fed with D3 food had the best albumin level (3.29±0.04 g/dl), which was substantially higher than the T_0 control (1.66±0.01 g/dl) (p < 0.05). The albumin level at the withdrawal period varied between 0.85±0.01 to 0.99±0.04. The value of albumin varied significantly (p < 0.005) between T₀, T₁, T₂ and T₃ in 0th day and pre challenge as well as between pre challenge and post challenge group fed with experimental diets. The results support the view of Talpur et al. (2012) [16] who reported an elevation in albumin level (p < 0.05) as compared to control when Asian sea bass was fed with garlic supplemented diets and post challenged with Virio harrveyi. Fish given Euglena powder showed a substantial change in albumin content at several assays between treatment groups and the control group, according to Das et al. (2009)^[3]. Same results were observed when Achyranthes aspera seed were added in the diet of Labeo rohita fingerlings.

Globulin

The profile of globulin in different treatment groups has been presented in (figure3). The globulin level of Amur carp at 0thday varied between 1.29±0.116 to 1.36±0.006 g/dl. The globulin level of Amur carp in pre challenged fishes varied between 1.63±0.02 to 1.79±0.05 g/dl among treatment. In pre challenged group, the maximum value for albumin was recorded as 1.79±0.05 g/dl (T₃) followed by 1.76±0.03 g/dl (T₂) and lowest in 1.63±0.02 g/dl (T₀). In case of post challenged fishes, the value ranged between 3.28±0.05 to 3.63±0.04 g/dl. The best level of globulin was found in treatment T₃ (3.63±0.04 g/dl). Fish blood biochemical examination revealed that T₃/D3, supplemented with 2.5g of each plant and 2.5g of chitosan, had the highest globulin level. Thus, treatment T₃ showed best globulin level which was significantly (p < 0.05) high as compared to control T₀ as well as other treatments. The globulin level at the withdrawal period varied between 1.66±0.02 to 1.82±0.05. The value of globulin varied significantly (p < 0.005)between T_0 , T_1 , T_2 and T_3 in 0^{th} day and pre challenge as well as between pre challenge and post challenge group fed with experimental diet. Similar results were reported by Harikrishnan et al., (2012)^[7] who reported elevation in globulin level in Epinephelus bruneus fed with 2% chitin or 1% chitosan diets. Subeena and Navarai (2012)^[15] reported enhanced (p < 0.05) serum protein, albumin, globulin content in Mystus keletius fed with herbal diets (Solanum trilobatum and Ocimum sanctum) as compared to the control group. Sahu et al., (2007)^[12] reported that Labeo rohita challenged with Aeromonas hydrophila showed significantly (p < 0.05)higher globulin level in all the groups in comparison with control when Curcuma longa was added in the diet. After feeding Sparus aurata with levamisole slight increment in globulin level has been reported by Mulero et al. (1998)^[10]. The globulin is certainly important for maintaining a good and healthy immune system (Jha et al., 2007)^[8].

A/G Ratio

The observations on A/G ratio in different treatment groups have been depicted in (figure 4). The A/G ratio of Amur carp at 0thday varied between 0.45 ± 0.006 to 0.47 ± 0.007 . The A/G ratio of Amur carp in pre challenge fishes varied between 0.50 ± 0.007 to 0.55 ± 0.006 among treatments. In pre challenged group, the maximum value for A/G ratio was recorded as 0.55 ± 0.006 (T₃) which was significantly higher from T_0 , T_1 and T_2 . In case of post challenged fishes, the value ranged between 0.50±0.007 to 0.90±0.009. Highest A/G ratio was obtained in T_3 (0.90±0.009). The value of A/G ratio varied significantly (p < 0.005) between T₀, T₁, T₂ and T₃ in 0th day and pre challenge as well as between pre challenge and post challenge group fed with the experimental diet. The value of A/G ratio during the withdrawal period ranged between 0.51 ± 0.006 to 0.54 ± 0.06 . Our results support the findings of Sivaram et al. (2004)^[13] who reported that the methanolic extracts of herbal plants like O. sanctum, W. somnifera and Myristica fragrans significantly improved the albumin–globulin (A/G) ratio in Epinephelus tauvina challenged with Vibrio harveyi. Chesti et al. (2018)^[2] reported that feeding of Allium sativum upto 1.5% in feed to fingerlings of Amur carp, Cyprinus carpio haematopterus, resulted in the significant changes (p < 0.05)between the pre challenge and post challenge groups in total serum protein, albumin, globulin, A/G ratio, aspartate aminotransferase (AST), serum alanine aminotransferase (ALT).

Alanine Aminotransferase (ALT)

The observations on alanine aminotransferase (ALT) in different treatment groups have been depicted in (figure5). The ALT of Amur carp at 0th day varied between 28.16±0.41 to 30.85±0.52. The ALT content in pre challenged fishes was in the range of 18.96 ± 0.501 (T₃) and 21.75 ± 0.386 (T₀). In case of post challenged fishes, the ALT content was recorded as 15.12±0.388 (T₃) to 19.63±0.314 (T₀). The value of ALT content among 0^{th} day and pre challenged as well as between pre challenged and post challenged fishes varied significantly (p < 0.05) between the treatments. The experiment conducted by Fazlolahzadeh et al. (2011)^[6] on Rainbow trout, Oncorynchu smykiss fed with garlic rich diet has resulted in variable values of ALT and their was no significant difference in treatments group as compared to control. The values of ALT after withdrawal period varied between 17.86±0.411 to 20.56±0.501. Increases in fish total serum protein, albumin, and globulin levels are hypothesized to be linked to a more robust innate immune response. An increase in the total leukocyte count, which is a significant source of serum protein production, including lysozyme, complement factors, and bactericidal peptides, could be the cause of the elevated serum protein content (Misra et al., 2006)^[9].



Fig 1: Total serum protein (g/dl) in different treatment groups of *Cyprinus carpio haematopterus* fed with different diet



Fig 2: Albumin content (g/dl) in different treatment groups of *Cyprinus carpio haematopterus* raised with different diet



Fig 3: Globulin content (g/dl) in different treatment groups of *Cyprinus carpio haematopterus* fed with different diets







Fig 5: Alanine Aminotransferase (IU/L) in different treatment groups of *Cyprinus carpio haematopterus* raised with different diets

Summary and Conclusion

The fish in this experiment fed diet D3 showed an increase in total serum protein, albumin, and globulin levels which may be an indication that the supplements incorporated in the diet had resulted in the enhancement of the non-specific immune response of fishes and also the increased levels of albumin may facilitate the transport of more humoral compounds as well as phytobiotic extract along with chitosan in the blood.

References

- Abdel-Tawwab M, Fayza Abbass E. Turmeric Powder, *Curcuma longa* L., in Common Carp, *Cyprinus carpio* L., Diets: Growth Performance, Innate Immunity, and Challenge against Pathogenic Aeromonas hydrophila Infection. Journal of theWorld aquaculture society. 2010;26:112-120.
- Chesti A, Chauhan RS, Khati A. Evaluation of immunostimulatory effect of dietary garlic (*Allium* sativum) in fingerlings of Amur carp, Cyprinus carpio haematopterus. The Pharma Innovation Journal. 2018;7(5):375-379
- Das BK, Pradhan J, Sahu S. The effect of Euglena viridis on immune response of rohu, *Labeo rohita* (Ham.) Fish and Shellfish Immunology. 2009;26:871-876.
- 4. The State of World Fisheries and Aquaculture 2015; FAO, Rome.
- 5. The State of World Fisheries and Aquaculture 2016; FAO, Rome.
- Fazlolahzadeh F, Keramati K, Nazifi S, Shirian S, Seifi S. Effect of Garlic (*Allium sativum*) on Hematological Parameters and Plasma Activities of ALT and AST of Rainbow trout in Temperature Stress. Australian Journal of Basic and Applied Sciences. 2011;5(9):84-90.
- Harikrishnan R, Kim Ju-Sang, Balasundaram C, Heo Moon-Soo. Immunomodulatory effects of chitin and chitosan enriched diets in *Epinephelus bruneus* against Vibrio alginolyticus infection. Aquaculture. 2012;326-329:46-52.
- Jha AK, Pal AK, Sahu NP, Kumar S, Mukherjee SC. Haemato-immunological responses to dietary yeast RNA, w-3 fatty acid and b-carotene in *Catla catla* juveniles. Fish Shellfish Immunology. 2007;23(5):17-27.
- Misra CK, Das BK, Mukherjee SC, Meher K. The Immunomodulatory effects of tufts in on the nonspecific immune system of Indian Major Carp Labeo rohita. Fish shellfish Immunology. 2006;20:728-738.
- Mulero V, Esteban MA, Munoz J, Mesegue J. Dietary intake of Levamisole enhances the immune response and disease resistance of marine teleost gilt head Sea bream (*Sparus aurata* L.). Fish and Shellfish Immunology. 1998;8:49-62.
- 11. Praveen Choyal, Dr. OP Garhwal, Dr. MR Choudhary, Dr. LN Bairwa, Dr. AC Shivran, Dr. DK Gothwal. Effect of sulphur and micronutrients on growth attributes of garlic (*Allium sativum* L.). Int. J Hortic Food Sci. 2022;4(2):223-226.
- 12. Sahu S, Das BK, Pradhan J, Mohapatra BC, Mishra BK, Sarangi N. Effect of *Magnifera indica* kernel as a feed additive on immunity and resistance to Aeromonas

hydrophila in Labeo rohita fingerlings. Fish Shellfish Immunology. 2007;23:109-118.

- 13. Sivaram V, Babu MM, Citarasu T, Immanuel G, Murugadass S, Marian MP. Growth and immuneresponse of juvenile greasy groupers (*Epinephelus tauvina*) fed with herbal antibacterial active principle supplemented diets against Vibrio harveyi infections. Aquaculture. 2004;237:9-20.
- Siwicki AK. Immunodulating influence of levamisole on non-specific immunity in carp (*Cyprinus carpio*). Development and Comparative Immunology. 1989;13:87-91.
- 15. Begum S, Navaraj PS. Synergistic effect of Plant Extracts Supplemented Diets on Immunity and Resistance to *Aeromonas hydrophila* in *Mystus keletius* Journal of Pharmacy and Biological Sciences. 2012;2(4):30-36.
- 16. Talpur AD, Mhd I. Dietary effects of garlic (*Allium sativum*) on haemato-immunological parameters, survival, growth, and disease resistance against *Vibrio harveyi* infection in Asian sea bass, Lates calcarifer (Bloch) Aquaculture. 2012;6:364-365.