

## International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693  
 ISSN Online: 2617-4707  
 IJABR 2024; SP-8(3): 631-633  
[www.biochemjournal.com](http://www.biochemjournal.com)  
 Received: 02-01-2024  
 Accepted: 13-02-2024

**Vishnoo Omar**  
 Assistant Professor,  
 Department of Entomology,  
 AKS University, Satna,  
 Madhya Pradesh, India

**Anand Shrivastava**  
 Assistant Professor,  
 Department of Entomology,  
 CSAUAT, Kanpur, Uttar  
 Pradesh, India

**Aditya Kumar**  
 Assistant Professor,  
 Department of Entomology,  
 AKS University, Satna,  
 Madhya Pradesh, India

**Amit Bagri**  
 Assistant Professor,  
 Department of Entomology,  
 AKS University, Satna,  
 Madhya Pradesh, India

**Vipin Kumar**  
 Assistant Professor,  
 Department of Extension,  
 AKS University, Satna,  
 Madhya Pradesh, India

**Pradip Kumar Patel**  
 SMS Agriculture, PANI  
 Sansthan, Balrampur, Uttar  
 Pradesh, India

**Arvind Kumar**  
 Assistant Professor,  
 Department of Entomology,  
 Haridwar University, Roorkee,  
 Uttarakhand, India

**Corresponding Author:**  
**Vishnoo Omar**  
 Assistant Professor,  
 Department of Entomology,  
 AKS University, Satna,  
 Madhya Pradesh, India

## Evaluate the effects of seed protectants on adult emergence of *C. chinensis*

**Vishnoo Omar, Anand Shrivastava, Aditya Kumar, Amit Bagri, Vipin Kumar, Pradip Kumar Patel and Arvind Kumar**

DOI: <https://doi.org/10.33545/26174693.2024.v8.i3Sh.849>

### Abstract

Chickpea (*Cicer arietinum* L.) is a richest source of protein, among various pulses to human diet and occupies an important position in the agriculture economy of our country. *Callosobruchus chinensis* L. is the major insect-pest and causes huge loss in storage besides affecting the quality by contaminating and decreasing the nutritive value of the grain. Study of the present investigation was carried out on growth and development of *C. chinensis* on Bengal gram (variety-Radhey) and to evaluate effects of seed protectants on adult emergence cent of *C. chinensis*. The least pupation was recorded. The neem leaf powder @ 15g/kg of seed and mustard oil @ 3ml/kg of seed gave lesser i.e. 1.36 and 1.86 percent adults emergence, which was at par i.e., 3.13.

**Keywords:** Chickpea (*Cicer arietinum* L.), seed, adult emergence

### Introduction

Pulses have been considered as the poor man's-meat which play an important role in food categories that have been extensively used as staple foods. The pulses contain about 20-25 percent protein which is higher than that of cereals. Hence, pulses are major source of protein to meet the increasing dietary demand in Indian people. The global pulse market is estimated about 60 million tonnes. For various reasons pulses are playing an important role in Indian agriculture. Chickpea is used in wide range of different preparation in our country and has a good source of energy 416 calories per 100g chicken along with protein 18 to 20 percent carbohydrate 51 to 70 percent 4 to 10 percent minerals, calcium, phosphorus and iron and vitamins also. In India, chickpea is the premier *rabi* followed by lentil, field pea and lathyrus. As far as production is concerned, India ranks first in the world and contributes around 70 percent (Anonymous, 2015) [2]. In Indian context, total pulse grown area is about 25.25 mh and production is only 16.47 mt (Directorate of economics and statistics, 2016). One of the major pulses cultivated and consumed in India, is chickpea and commonly known as Gram, Bengal gram or White gram. India is the major producing country for chickpea, contributing for over 75% of total production in the world (Anonymous, 2016) [3]. *Callosobruchus chinensis* L. is cosmopolitan and a serious pest of pulses and has also been reported attacking cotton seed, sorghum and maize (Ahmed *et al.*, 2003) [1]. Due to invasion of the beetle, deterioration in quality and quantity of stored pulse. The grubs bore into the pulse grains which become unsuitable for human consumption and for the production of sprouts. This is important pests of pulse crop in India under storage conditions (Tapondjou *et al.*, 2002) [15] and cause annual loss of nearly 0.21 million tonnes amounting to rupees 315 million. The extensive application of insecticides is directly related with the development of resistance of stored product insect species, raising serious concerns for human health as a result of food contamination with residues and possible environmental hazards (Boyer *et al.* 2012; Stadler *et al.* 2012) [4, 13]. Serious problems of genetic resistance by insect species, pest resurgence, residual toxicity, phytotoxicity, vertebrate toxicity, widespread environmental hazards and increasing cost of application of the presently used synthetic pesticides have directed the need for effective biodegradable pesticides (Talukdar and Howse, 1994) [14]. This awareness has created worldwide interest in the development of alternative strategic including the re-examination of using plant derivative materials is more readily biodegradable.

Some are less toxic to mammals, may be more selective in action and may disgrace the development of resistance. The main advantage is that they may be easily and cheaply produced by farmers and small-scale industries as crude or partially purified extracts. In two decades, Considerable efforts have been made at screening plants in order to develop new botanical insecticides as alternative to the existing insecticide. The present investigation was carried out on growth and development of *C. chinensis* on Bengal gram (variety-Radhey) and to evaluate effects of seed protectants on adult emergence of *C. chinensis*.

### Materials and Methods

The present investigation was carried out on growth and development of *C. chinensis* on Bengal gram (variety-Radhey) and to evaluate effects of seed protectants adult emergence of *C. chinensis*.

**Test insect:** *Callosobruchus chinensis* Linn.

### Taxonomy of insect

Phylum	:	Arthropoda
Class	:	Insecta
Order	:	Coleoptera
Supper Family	:	Chrysomelidea
Family	:	Bruchidae
Genus	:	<i>Callosobruchus</i>
Species	:	<i>chinensis</i>

The present study was conducted in the Insectary, Department of Agricultural Entomology, Chandra Shekhar Azad University of agriculture and technology, Kanpur, Madurai during 2017-18. The chickpea variety *Radhey* was selected to conduct the investigation on *C. chinensis* which was obtained from Economic Botanist (legumes), C. S. A. University of Agriculture and Technology Kanpur. The adults of pulse beetle were initially collected from infested stock of local granaries and brought to the laboratory for the mass culture and rearing on gram variety (Radhey) kept in 2 kg capacity plastic container filled with 1 kg chickpea. 20 freshly emerged pairs of males and females of pulse beetle were released in a plastic container filled with fresh seeds of

gram and covered with muslin cloth and tied around with rubber band and kept at room temperature. The food of container culture was replaced with fresh one, as and when it is needed. The newly emerged adult from the culture were used in the present study. The seeds of gram were properly sterilized in an electric oven at a temperature of 55<sup>o</sup>c for 24 hours to make it free from any contamination. The seeds were again conditioned in regulated temperature and 75 ± 5% relative humidity under desiccators. The experiments were based on “complete randomized design” (CRD) with 7 treatments and 3 replications. The 50 g grains of gram was weighted with the help of physical balance. Five pairs of *C. chinensis* of same age obtained from the laboratory culture were introduced in each jar, covered by muslin cloth and tied with rubber band and kept under the room temperature in the P.G. laboratory of the department. With a view to study the adult emergence.

### Results and Discussion

The data depicted in (Table 1) showed that all the grain protectants were found at par effective which provided reduced adults emergence. The neem leaf powder @ 15g/kg of seed and mustard oil @ 3ml/kg of seed gave lesser i.e. 1.36 and 1.86 percent adults emergence, respectively, followed by dhatura leaf powder @ 15g/kg of seed, linseed oil @ 3 ml/kg of seed, castor leaf powder @ 15g/kg of seed, coconut oil @ 3ml/kg of seed and oak leaf powder @ 15g/kg of seed which provided 2.18, 2.24, 2.80, 3.05, and 3.13 percent reduced adults emergence, respectively, as compared to 67.14 percent reduced adults emergence in control.

These findings corroborate with those of Chudasama *et al.* (2015)<sup>[5]</sup> on *C. maculatus* with 5% black pepper seed extract and neem seed kernel extract. The effect of *O. sanctum* inhibited adult emergence of *C. maculatus* (Ratnasekera and Rajapakse, 2010)<sup>[8]</sup>. also found that neem leaf powder @ 15 g/kg giving protection to cowpea against *C. chinensis*. These results are in accordance with the finding of Pandey *et al.* (1986)<sup>[6]</sup> who found that neem leaves and twigs extracts @ 0.5, 1.0 and 1.5 parts /100 parts of (w/w) green gram seeds were highly repellent to the *C. chinensis* and protecting against the infestation.

**Table:** Effect of different botanicals on adult's emergence percent of *C. chinensis* L. in chickpea grains

Sr. No.	Treatment	Dosage (g/ml) per kg of seed	Adults Emergence percent
1.	Neem Leaf Powder	15g	1.36 (1.36)
2.	Mustard Oil	3ml	1.86 (1.53)
3.	Dhatura Leaf Powder	15g	2.18 (1.63)
4.	Linseed Oil	3ml	2.24 (1.65)
5.	Castor Leaf Powder	15g	2.80 (1.81)
6.	Coconut Oil	3ml	3.05 (1.88)
7.	Oak Leaf Powder	15g	3.13 (1.90)
8.	Control		67.14 (8.22)
	CD 5%		1.93
	SE, (d)		0.91

### Conclusion

On the basis of results which was obtained present experiment, the following conclusion are brought out, which may be useful for scientists, research workers and farmers.

The neem leaf powder @ 15 g/kg of seed and mustard oil @ 3ml/kg of seed gave lesser i.e. 1.36 and 1.86 percent adults emergence, respectively, followed by dhatura leaf powder @ 15 g/kg of seed, linseed oil @ 3 ml/kg of seed, castor leaf powder @ 15 g/kg of seed, coconut oil @ 3ml/kg of seed

and oak leaf powder @ 15g/kg of seed which provided 2.18, 2.24, 2.80, 3.05, and 3.13 percent reduced. Adults emergence, respectively, as compared to 67.14 percent reduced adults emergence in control.

### References

- Ahmed K, Itino T, Ichikawa T. Duration of developmental stages of *Callosobruchus chinensis* (Coleoptera: Bruchidae) on Adzuki bean and the effects

- of *neem* and sesame oils at different stages of their development, 2003.
2. Anonymous. Project Coordinator's Report. AICRP on Chickpea, IIPR, Kanpur, 2015, 36.
  3. Anonymous. Project Coordinator's Report. AICRP on Chickpea, IIPR, Kanpur, 2016.
  4. Boyer S, Zhang H, Lemperiori G. A review of control methods and resistance mechanisms in stored-product insects. *Bulletin of entomological research* 2012;102(2):213-229.
  5. Chudasama JA, Sagarka NB, Sharma S. Deterrent effect of plant extracts against *Callosobruchus maculatus* on stored cowpea in Saurashtra (Gujarat, India). *Journal of Applied and Natural Science*. 2015;7(1):187-191.
  6. Pandey S, Mathur ND CK, Tripathi RA. Effect of protectants against pulse beetle *Callosobruchus chinensis* L. *Journal of Entomology*. 1986;48(1):85-90.
  7. Pandey NK, Singh SG. Effect of *neem* bark powder infestation of pulse beetle *Callosobruchus chinensis*. *Indian Journal of Entomology*. 1997;59(2):161-163.
  8. Ratnasekera D, Rajapakse R. Repellent properties of plant oil vapours on pulse beetle (*Callosobruchus maculatus* L.) (Coleoptera: Bruchidae) in stored green gram (*Vigna radiata* L.). *Tropical Agricultural Research and Extension*. 2010;12(1):26-31.
  9. Singal SK. Biology of pulse beetle, *Callosobruchus chinensis* (L.) on cowpea under field and laboratory conditions. *Journal of Insect Science*. 1998;11(2):130-132.
  10. Singh R. Evaluation of some plant products for their oviposition deterrent properties against the *Callosobruchus maculatus* (F.) on chick pea seeds. *Journal of Agriculture & Technology*. 2011;7:1363-1367.
  11. Singh S, Odak SC, Singh Z. Studies on preference of pulse beetle, *Callosobruchus chinensis* L. for different hosts. *Bulletin of Grain Technology*. 1977;15(1):20-26.
  12. Singh SC. Effect of neem bark powder as protectant against *Callosobruchus chinensis* L. on stored chickpea and lentil. *Indian Journal of Entomology*. 2005;67(3):203-205.
  13. Stadler T, Buteler M, Weaver DK, Sofie S. Comparative toxicity of nanostructured alumina and a commercial inert dust for *Sitophilus oryzae* (L.) and *Rhyzopertha dominica* (F.) at varying ambient humidity levels. *Journal of stored products research*. 2012;48:81-90.
  14. Talukdar FA, Howse PE. Repellent toxic and food protectant effects of *pithraj*, *Aphanamixis polystachya* (Meliaceae) against pulse beetle, *Callosobruchus chinensis* (L.) in storage. *Journal of Chemical Ecology*. 1994;20(4):899-908.
  15. Tapondjou LA, Adler C, Bouda H, Fontem DA. Efficacy of powder and essential oil from *Chenopodium ambrosioides* leaves as post-harvest grain protectants against six stored product beetles. *Journal of Stored Products and Research*. 2002;38:395-402.
  16. Tiroesele B, Thomas K, Seketeme S. Control of cowpea weevil, *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae), using natural plant products. *Monthly Insects*. 2015;6(1):77-84.